

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

A183

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup>:</b> <b>C07H 21/04, C07K 14/705, C12N 15/09, 15/63, C12Q 1/68</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 98/54202</b> <b>(43) International Publication Date:</b> 3 December 1998 (03.12.98)
<b>(21) International Application Number:</b> PCT/US98/10981 <b>(22) International Filing Date:</b> 29 May 1998 (29.05.98)  <b>(30) Priority Data:</b> 60/050,936 30 May 1997 (30.05.97) US 60/069,112 9 December 1997 (09.12.97) US  <b>(71) Applicant (for all designated States except US):</b> HUMAN GENOME SCIENCES, INC. [US/US]; 9410 Key West Avenue, Rockville, MD 20850 (US).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> NI, Jian [CN/US]; 5502 Manorfield Road, Rockville, MD 20853 (US). ROSEN, Craig, A. [US/US]; 22400 Rolling Hill Road, Laytonsville, MD 20882 (US).  <b>(74) Agents:</b> BROOKES, A., Anders et al.; Human Genome Sciences, Inc., 9410 Key West Avenue, Rockville, MD 20850 (US).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> <i>With an indication in relation to deposited biological material furnished under Rule 13bis separately from the description.</i>
<b>(54) Title:</b> HUMAN TUMOR NECROSIS FACTOR RECEPTOR TR10  <b>(57) Abstract</b>  The present invention relates to a novel protein, TR10, which is a member of the tumor necrosis factor (TNF) receptor superfamily and the TRAIL receptor subfamily. In particular, isolated nucleic acid molecules are provided encoding the human TR10 protein. TR10 polypeptides are also provided as are vectors, host cells and recombinant methods for producing the same. The invention further relates to screening methods for identifying agonists and antagonists of TR10 activity.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## Human Tumor Necrosis Factor Receptor TR10

### *Background of the Invention*

#### *Field of the Invention*

5       The present invention relates to a novel member of the tumor necrosis factor family of receptors. More specifically, isolated nucleic acid molecules are provided encoding a novel human tumor necrosis factor receptor, TR10. TR10 polypeptides are also provided, as are vectors, host cells, and recombinant methods for producing the same. The invention further relates to screening methods for identifying agonists and  
10       antagonists of TR10 activity.

#### *Related Art*

      Many biological actions, for instance, response to certain stimuli and natural biological processes, are controlled by factors, such as cytokines. Many cytokines act  
15       through receptors by engaging the receptor and producing an intra-cellular response.

      For example, tumor necrosis factors (TNF) alpha and beta are cytokines, which act through TNF receptors to regulate numerous biological processes, including protection against infection and induction of shock and inflammatory disease. The TNF molecules belong to the "TNF-ligand" superfamily, and act together with their receptors or counter-ligands, the "TNF-receptor" superfamily. So far, nine members of the TNF  
20       ligand superfamily have been identified and ten members of the TNF-receptor superfamily have been characterized.

      Among the ligands there are included TNF- , lymphotoxin- (LT- , also known as TNF- ), LT- (found in complex heterotrimer LT- 2- ), FasL, CD40L, CD27L, CD30L, 4-1BBL, OX40L and nerve growth factor (NGF). The superfamily of  
25       TNF receptors includes the p55TNF receptor, p75TNF receptor, TNF receptor-related protein, FAS antigen or APO-1, CD40, CD27, CD30, 4-1BB, OX40, low affinity p75 and NGF-receptor (A. Meager, *Biologicals* 22:291-295 (1994)).

      Many members of the TNF-ligand superfamily are expressed by activated T-  
30       cells, implying that they are necessary for T-cell interactions with other cell types which underlie cell ontogeny and functions. (A. Meager, *supra*).

      Considerable insight into the essential functions of several members of the TNF receptor family has been gained from the identification and creation of mutants that abolish the expression of these proteins. For example, naturally occurring mutations in  
35       the FAS antigen and its ligand cause lymphoproliferative disease (R. Watanabe-Fukunaga *et al.*, *Nature* 356:314 (1992)), perhaps reflecting a failure of programmed

cell death. Mutations of the CD40 ligand cause an X-linked immunodeficiency state characterized by high levels of immunoglobulin M and low levels of immunoglobulin G in plasma, indicating faulty T-cell-dependent B-cell activation (R.C. Allen *et al.*, *Science* 259:990 (1993)). Targeted mutations of the low affinity nerve growth factor  
5 receptor cause a disorder characterized by faulty sensory innervation of peripheral structures (K.F. Lee *et al.*, *Cell* 69:737 (1992)).

TNF and LT- are capable of binding to two TNF receptors (the 55- and 75-kd TNF receptors). A large number of biological effects elicited by TNF and LT- , acting through their receptors, include hemorrhagic necrosis of transplanted tumors,  
10 cytotoxicity, a role in endotoxic shock, inflammation, immunoregulation, proliferation and anti-viral responses, as well as protection against the deleterious effects of ionizing radiation. TNF and LT- are involved in the pathogenesis of a wide range of diseases, including endotoxic shock, cerebral malaria, tumors, autoimmune disease, AIDS and graft-host rejection (B. Beutler and C. Von Huffel, *Science* 264:667-668 (1994)).  
15 Mutations in the p55 receptor cause increased susceptibility to microbial infection.

Moreover, an about 80 amino acid domain near the C-terminus of TNFR1 (p55) and Fas was reported as the "death domain," which is responsible for transducing signals for programmed cell death (Tartaglia *et al.*, *Cell* 74:845 (1993)).

Apoptosis, or programmed cell death, is a physiologic process essential to the  
20 normal development and homeostasis of multicellular organisms (H. Steller, *Science* 267:1445-1449 (1995)). Derangements of apoptosis contribute to the pathogenesis of several human diseases including cancer, neurodegenerative disorders, and acquired immune deficiency syndrome (C.B. Thompson, *Science* 267:1456-1462 (1995)). Recently, much attention has focused on the signal transduction and biological function  
25 of two cell surface death receptors, Fas/APO-1 and TNFR-1 (J.L. Cleveland *et al.*, *Cell* 81:479-482 (1995); A. Fraser *et al.*, *Cell* 85:781-784 (1996); S. Nagata *et al.*, *Science* 267:1449-56 (1995)). Both are members of the TNF receptor family, which also include TNFR-2, low affinity NGFR, CD40, and CD30, among others (C.A. Smith *et al.*, *Science* 248: 1019-23 (1990); M. Tewari *et al.*, in *Modular Texts in Molecular and*  
30 *Cell Biology* M. Purton, Heldin, Carl, Ed. (Chapman and Hall, London, 1995). While family members are defined by the presence of cysteine-rich repeats in their extracellular domains, Fas/APO-1 and TNFR-1 also share a region of intracellular homology, appropriately designated the "death domain," which is distantly related to the *Drosophila* suicide gene, reaper (P. Golstein *et al.*, *Cell* 81:185-6 (1995); K. White *et al.*, *Science* 264:677-83 (1994)). This shared death domain suggests that both  
35 receptors interact with a related set of signal transducing molecules that, until recently,

remained unidentified. Activation of Fas/APO-1 recruits the death domain-containing adapter molecule FADD/MORT1 (A.M. Chinnaiyan *et al.*, *Cell* 81:505-512 (1995); M. P. Boldin *et al.*, *J. Biol. Chem.* 270:7795-8 (1995); F.C. Kischkel *et al.*, *EMBO* 14:5579-5588 (1995)), which in turn binds and presumably activates FLICE/MACH1, a member of the ICE/CED-3 family of pro-apoptotic proteases (M. Muzio *et al.*, *Cell* 85: 817-827 (1996); M.P. Boldin *et al.*, *Cell* 85:803-815 (1996)). While the central role of Fas/APO-1 is to trigger cell death, TNFR-1 can signal an array of diverse biological activities-many of which stem from its ability to activate NF-kB (L.A. Tartaglia *et al.*, *Immunol Today* 13:151-153 (1992)). Accordingly, TNFR-1 recruits the multivalent adapter molecule TRADD, which like FADD, also contains a death domain (H. Hsu *et al.*, *Cell* 81:495-504 (1995); H. Hsu *et al.*, *Cell* 84:299-308 (1996)). Through its associations with a number of signaling molecules including FADD, TRAF2, and RIP, TRADD can signal both apoptosis and NF-kB activation (H. Hsu *et al.*, *Cell* 84:299-308 (1996); H. Hsu *et al.*, *Immunity* 4:387-396 (1996)).

Recently, a new apoptosis inducing TNF ligand has been discovered. S.R. Wiley *et al.*, *Immunity* 3:673-682 (1995), named the new molecule, "TNF-related apoptosis-inducing ligand" or "TRAIL." R.M. Pitti *et al.*, *J. Biol. Chem.* 271:12687-12690 (1996), named the molecule "Apo-2 ligand" or "Apo-2L." This molecule was also disclosed in co-pending U.S. provisional patent application no. 60/013405. For convenience, this molecule will be referred to herein as TRAIL.

Unlike FAS ligand, whose transcripts appear to be largely restricted to stimulated T-cells, significant levels of TRAIL are detected in many human tissues (e.g., spleen, lung, prostate, thymus, ovary, small intestine, colon, peripheral blood lymphocytes, placenta, kidney), and it is constitutively transcribed by some cell lines. It has been shown that TRAIL acts independently from the FAS ligand (S.R. Wiley *et al.*, *supra*). It has also been shown that TRAIL activates apoptosis rapidly, within a time frame that is similar to death signaling by Fas/Apo-1L, but much faster than TNF-induced apoptosis. S.A. Marsters *et al.*, *Current Biology* 6:750-752 (1996). The inability of TRAIL to bind TNFR-1, Fas, or the recently identified DR3, suggests that TRAIL may interact with a unique receptor(s).

Work to date suggests that there are several unique TNF receptors for TRAIL. In co-pending U.S. provisional patent application no. 60/035,722, one novel death domain containing receptor for TRAIL, DR4, was disclosed. See, Pan *et al.*, *Science* 276:111-113 (April 1997). In co-pending U.S. provisional patent application no. 60/040,846, a novel death domain containing receptor, DR5 (TR7), was disclosed. This receptor has now been shown to bind TRAIL. In co-pending U.S. provisional

patent application no. 60/035,496, another receptor, TR5, was disclosed. This receptor has also now been shown to bind TRAIL, however, TR5 has been shown to be a non-signaling decoy receptor which antagonizes apoptosis.

The effects of TNF family ligands and receptors are varied and influence numerous functions, both normal and abnormal, in the biological processes of the mammalian system. There is a clear need, therefore, for identification and characterization of such receptors and ligands that influence biological activity, both normally and in disease states. In particular, there is a need to isolate and characterize additional novel receptors that bind TRAIL.

### *Summary of the Invention*

The present invention provides isolated nucleic acid molecules comprising a polynucleotide encoding the TR10 receptor having the amino acid sequence shown in SEQ ID NO:2 or the amino acid sequence encoded by the cDNA clone deposited as American Type Culture Collection ("ATCC") Deposit No. 209040 on May 15, 1997. The ATCC is located at 10801 University Boulevard, Manassas, Virginia 20110-2209.

The present invention also relates to recombinant vectors, which include the isolated nucleic acid molecules of the present invention, and to host cells containing the recombinant vectors, as well as to methods of making such vectors and host cells and for using them for production of TR10 polypeptides or peptides by recombinant techniques.

The invention further provides an isolated TR10 polypeptide having an amino acid sequence encoded by a polynucleotide described herein.

The present invention also provides diagnostic assays such as quantitative and diagnostic assays for detecting levels of TR10 protein. Thus, for instance, a diagnostic assay in accordance with the invention for detecting over-expression of TR10, or soluble form thereof, compared to normal control tissue samples may be used to detect the presence of tumors.

Tumor Necrosis Factor (TNF) family ligands are known to be among the most pleiotropic cytokines, inducing a large number of cellular responses, including cytotoxicity, anti-viral activity, immunoregulatory activities, and the transcriptional regulation of several genes. Cellular response to TNF-family ligands include not only normal physiological responses, but also diseases associated with increased apoptosis or the inhibition of apoptosis. Apoptosis-programmed cell death-is a physiological mechanism involved in the deletion of peripheral T lymphocytes of the immune system, and its dysregulation can lead to a number of different pathogenic processes. Diseases associated with increased cell survival, or the inhibition of apoptosis, include cancers,



autoimmune disorders, viral infections, inflammation, graft vs. host disease, acute graft rejection, and chronic graft rejection. Diseases associated with increased apoptosis include AIDS, neurodegenerative disorders, myelodysplastic syndromes, ischemic injury, toxin-induced liver disease, septic shock, cachexia, and anorexia.

5        Thus, the invention further provides a method for inhibiting apoptosis induced by a TNF-family ligand, which involves administering to a cell which expresses the TR10 polypeptide an effective amount of an agonist capable of increasing TR10 mediated signaling. Preferably, TR10 mediated signaling is increased to treat a disease wherein increased apoptosis is exhibited.

10        In a further aspect, the present invention is directed to a method for enhancing apoptosis induced by a TNF-family ligand, which involves administering to a cell which expresses the TR10 polypeptide an effective amount of an antagonist capable of decreasing TR10 mediated activity. Preferably, TR10 mediated activity is decreased to treat a disease wherein decreased apoptosis is exhibited.

15        Whether any candidate "agonist" or "antagonist" of the present invention can enhance or inhibit apoptosis can be determined using art-known TNF-family ligand/receptor cellular response assays, including those described in more detail below. Thus, in a further aspect, a screening method is provided for determining whether a candidate agonist or antagonist is capable of enhancing or inhibiting a cellular  
20        response to a TNF-family ligand. The method involves contacting cells which express the TR10 polypeptide with a candidate compound and a TNF-family ligand, assaying a cellular response, and comparing the cellular response to a standard cellular response, the standard being assayed when contact is made with the ligand in absence of the candidate compound, whereby an increased cellular response over the standard  
25        indicates that the candidate compound is an agonist of the ligand/receptor signaling pathway and a decreased cellular response compared to the standard indicates that the candidate compound is an antagonist of the ligand/receptor signaling pathway. By the invention, a cell expressing the TR10 polypeptide can be contacted with either an endogenous or exogenously administered TNF-family ligand.

30

### ***Brief Description of the Figures***

Figure 1 shows the nucleotide (SEQ ID NO:1) and deduced amino acid sequence (SEQ ID NO:2) of the TR10 receptor. Predicted amino acids 1-55 constitute the signal peptide (amino acid residues from about -55 to about -1 in SEQ ID NO:2);  
35        amino acids 56-212 constitute the extracellular domain (amino acid residues from about 1 to about 157 in SEQ ID NO:2); amino acids 213-230 constitute the transmembrane domain (amino acid residues from about 158 to about 175 in SEQ ID NO:2); and amino

acids 231-386 constitute the intracellular domain (amino acid residues from about 176 to about 331 in SEQ ID NO:2), of which amino acids 353-363 constitute the partial death domain (amino acid residues from about 298 to about 308 in SEQ ID NO:2).

Figure 2 shows the regions of similarity between the amino acid sequences of the TR10 receptor protein (SEQ ID NO:2), and the Fas receptor (SEQ ID NO:3), NGFR p75 (SEQ ID NO:4), human TNFR 1 (SEQ ID NO:5), and DR4 (SEQ ID NO:6).

Figure 3 shows an analysis of the TR10 amino acid sequence. Alpha, beta, turn and coil regions; hydrophilicity and hydrophobicity; amphipathic regions; flexible regions; antigenic index and surface probability are shown. In the "Antigenic Index - Jameson-Wolf" graph, amino acid residues about 57 to about 113, about 130 to about 197, and about 250 to about 283 in Figure 1 correspond to the shown highly antigenic regions of the TR10 protein. These highly antigenic fragments in Figure 1 correspond to the following fragments, respectively, in SEQ ID NO:2: amino acid residues from about 2 to about 58, from about 75 to about 142, and from about 195 to about 228.

#### ***Detailed Description of the Preferred Embodiments***

The present invention provides isolated nucleic acid molecules comprising a polynucleotide encoding a TR10 polypeptide having the amino acid sequence shown in FIG. 1 (SEQ ID NO:2), which was determined by sequencing a cloned cDNA. The TR10 polypeptide of the present invention shares sequence homology with human NGFR, TNFRI, DR4, and Fas (FIG. 2). The nucleotide sequence shown in FIG. 1 (SEQ ID NO:1) was obtained by sequencing a cDNA clone, which was deposited on May 15, 1997 at the American Type Culture Collection, 12301 Park Lawn Drive, Rockville, Maryland 20852, and given Accession Number 209040. The deposited clone is inserted in the pCMVSPORT 2.0 plasmid (Life Technologies, Rockville, MD) using the Sal I/Not I restriction endonuclease cleavage sites.

#### ***Nucleic Acid Molecules***

Unless otherwise indicated, all nucleotide sequences determined by sequencing a DNA molecule herein were determined using an automated DNA sequencer (such as the Model 373 from Applied Biosystems, Inc.), and all amino acid sequences of polypeptides encoded by DNA molecules determined herein were predicted by translation of a DNA sequence determined as above. Therefore, as is known in the art for any DNA sequence determined by this automated approach, any nucleotide sequence determined herein may contain some errors. Nucleotide sequences determined by automation are typically at least about 90% identical, more typically at



least about 95% to at least about 99.9% identical to the actual nucleotide sequence of the sequenced DNA molecule. The actual sequence can be more precisely determined by other approaches including manual DNA sequencing methods well known in the art. As is also known in the art, a single insertion or deletion in a determined nucleotide sequence compared to the actual sequence will cause a frame shift in translation of the nucleotide sequence such that the predicted amino acid sequence encoded by a determined nucleotide sequence will be completely different from the amino acid sequence actually encoded by the sequenced DNA molecule, beginning at the point of such an insertion or deletion.

Using the information provided herein, such as the nucleic acid sequence set out in SEQ ID NO:1, a nucleic acid molecule of the present invention encoding a TR10 polypeptide may be obtained using standard cloning and screening procedures, such as those for cloning cDNAs using mRNA as starting material. Illustrative of the invention, the nucleic acid molecule described in SEQ ID NO:1 was discovered in a cDNA library derived from keratinocytes. The gene of the present invention has also been identified in cDNA libraries from the following tissues: fetal liver, peripheral blood lymphocytes (PBL), lung, kidney, small intestine, colon, endothelial cells, and monocyte activated tissue. Furthermore, the following cancer cell lines express TR10: Hela cell S3, SW480 (colorectal adenocarcinoma), and A549 (lung carcinoma).

The determined nucleotide sequence of the TR10 cDNA of SEQ ID NO:1 contains an open reading frame encoding a protein of about 331 amino acid residues, with a predicted leader sequence of about 55 amino acid residues, and a deduced molecular weight of about 42 kDa. The amino acid sequence of the predicted mature TR10 receptor is shown in SEQ ID NO:2 from amino acid residue about 1 to residue about 331. Of known members of the TNF receptor family, the TR10 polypeptide of the invention shares the greatest degree of homology with human DR4 (See FIG. 2), including significant sequence homology over multiple cysteine rich domains.

Owing to the sequence homology exhibited between TR10 and DR4 (and other death domain containing receptors), it was immediately recognized that TR10 would likely also bind to TRAIL. The cytoplasmic domain, interestingly, contains only a partial (or truncated) death domain. As described in Example 5, below, TR10 binds TRAIL but does not appear to cause cell death. TR10 binding of TRAIL, to the contrary, antagonizes apoptosis. Such antagonistic effect on TRAIL induced apoptosis can be achieved both through ectopic expression of TR10 and through exogenous administration of soluble TR10.

To examine the tissue distribution of TR10, Northern blot analysis was performed. A single transcript was detected in multiple human tissues at varying levels of expression, including, heart, lung, brain, placenta, liver, skeletal muscle, kidney, pancreas, spleen, thymus, prostate, testis, ovary, small intestine, colon, PBLs, lymph  
5 node, bone marrow and fetal liver. TR10 expression was not observed in most cancer cell lines tested. See Example 4, below.

As indicated, the present invention also provides the mature form(s) of the TR10 receptor of the present invention. According to the signal hypothesis, proteins secreted by mammalian cells have a signal or secretory leader sequence which is cleaved  
10 from the mature protein once export of the growing protein chain across the rough endoplasmic reticulum has been initiated. Most mammalian cells and even insect cells cleave secreted proteins with the same specificity. However, in some cases, cleavage of a secreted protein is not entirely uniform, which results in two or more mature species on the protein. Further, it has long been known that the cleavage specificity of  
15 a secreted protein is ultimately determined by the primary structure of the complete protein, that is, it is inherent in the amino acid sequence of the polypeptide.

Therefore, the present invention provides a nucleotide sequence encoding the mature TR10 polypeptide having the amino acid sequence encoded by the cDNA clone contained in the host identified as ATCC Deposit No. 209040, and as shown in Figure  
20 1 (SEQ ID NO:2). By the mature TR10 protein having the amino acid sequence encoded by the cDNA clones contained in the host identified as ATCC Deposit No. 209040 is meant the mature form(s) of the TR10 receptor produced by expression in a mammalian cell (e.g., COS cells, as described below) of the complete open reading frame encoded by the human DNA sequence of the clone contained in the vector in the  
25 deposited host. As indicated below, the mature TR10 receptor having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. 209040, may or may not differ from the predicted mature TR10 protein shown in SEQ ID NO:2 (amino acids from about 1 to about 331) depending on the accuracy of the predicted cleavage site based on computer analysis.

30 Methods for predicting whether a protein has a secretory leader as well as the cleavage point for that leader sequence are available. For instance, the method of McGeoch (*Virus Res.* 3:271-286 (1985)) and von Heinje (*Nucleic Acids Res.* 14:4683-4690 (1986)) can be used. The accuracy of predicting the cleavage points of known mammalian secretory proteins for each of these methods is in the range of 75-80%.  
35 von Heinje, *supra*. However, the two methods do not always produce the same predicted cleavage point(s) for a given protein.

In the present case, the predicted amino acid sequence of the complete TR10 polypeptide of the present invention was analyzed by a computer program ("PSORT"). See K. Nakai and M. Kanehisa, *Genomics* 14:897-911 (1992). PSORT is an expert system for predicting the cellular location of a protein based on the amino acid sequence. As part of this computational prediction of localization, the methods of McGeoch and von Heinje are incorporated. The analysis by the PSORT program predicted the cleavage site between amino acids -1 and 1 in SEQ ID NO:2. Thereafter, the complete amino acid sequences were further analyzed by visual inspection, applying a simple form of the (-1,-3) rule of von Heinje. von Heinje, *supra*. Thus, the leader sequence for the TR10 protein is predicted to consist of amino acid residues from about -55 to about -1 in SEQ ID NO:2, while the mature TR10 protein is predicted to consist of residues from about 1-331 in SEQ ID NO:2.

As one of ordinary skill would appreciate, due to the possibilities of sequencing errors, as well as the variability of cleavage sites for leaders in different known proteins, the predicted TR10 polypeptide encoded by the deposited cDNA comprises about 386 amino acids, but may be anywhere in the range of 376-396 amino acids; and the predicted leader sequence of this protein is about 55 amino acids, but may be anywhere in the range of about 45 to about 65 amino acids.

As indicated, nucleic acid molecules of the present invention may be in the form of RNA, such as mRNA, or in the form of DNA, including, for instance, cDNA and genomic DNA obtained by cloning or produced synthetically. The DNA may be double-stranded or single-stranded. Single-stranded DNA may be the coding strand, also known as the sense strand, or it may be the non-coding strand, also referred to as the anti-sense strand.

By "isolated" nucleic acid molecule(s) is intended a nucleic acid molecule, DNA or RNA, which has been removed from its native environment. For example, recombinant DNA molecules contained in a vector are considered isolated for the purposes of the present invention. Further examples of isolated DNA molecules include recombinant DNA molecules maintained in heterologous host cells or purified (partially or substantially) DNA molecules in solution. Isolated RNA molecules include *in vivo* or *in vitro* RNA transcripts of the DNA molecules of the present invention. Isolated nucleic acid molecules according to the present invention further include such molecules produced synthetically.

Isolated nucleic acid molecules of the present invention include DNA molecules comprising an open reading frame (ORF) shown in SEQ ID NO:1; DNA molecules comprising the coding sequence for the mature TR10 protein; and DNA molecules

which comprise a sequence substantially different from those described above, but which, due to the degeneracy of the genetic code, still encode the TR10 protein. Of course, the genetic code is well known in the art. Thus, it would be routine for one skilled in the art to generate such degenerate variants.

5 In addition, the invention provides nucleic acid molecules having nucleotide sequences related to extensive portions of SEQ ID NO:1 which have been determined from the following related cDNA clones: HSABD50R (SEQ ID NO:7), HGBDL20R(SEQ ID NO:8), and HELDL61R (SEQ ID NO:9), and AA150849 (SEQ ID NO:15).

10 In another aspect, the invention provides isolated nucleic acid molecules encoding the TR10 polypeptide having an amino acid sequence as encoded by the cDNA clone contained in the plasmid deposited as ATCC Deposit No. 209040 on May 15, 1997. In a further embodiment, nucleic acid molecules are provided that encode the mature TR10 polypeptide or the full length TR10 polypeptide lacking the N-terminal  
15 methionine. The invention further provides an isolated nucleic acid molecule having the nucleotide sequence shown in SEQ ID NO:1 or the nucleotide sequence of the TR10 cDNA contained in the above-described deposited clone, or a nucleic acid molecule having a sequence complementary to one of the above sequences. Such isolated molecules, particularly DNA molecules, are useful as probes for gene mapping by *in*  
20 *situ* hybridization with chromosomes, and for detecting expression of the TR10 gene in human tissue, for instance, by Northern blot analysis.

The present invention is further directed to fragments of the isolated nucleic acid molecules described herein. By a fragment of an isolated DNA molecule having the nucleotide sequence of the deposited cDNA or the nucleotide sequence shown in SEQ  
25 ID NO:1 is intended DNA fragments at least about 15nt, and more preferably at least 20 nt, still more preferably at least about 30 nt, and even more preferably, at least about 40 nt in length which are useful as diagnostic probes and primers as discussed herein. Of course, larger fragments 50-1500 nt in length are also useful according to the present invention, as are fragments corresponding to most, if not all, of the nucleotide sequence  
30 of the deposited cDNA or as shown in SEQ ID NO:1. By a fragment at least 20 nt in length, for example, is intended fragments which include 20 or more contiguous bases from the nucleotide sequence of the deposited cDNA or the nucleotide sequence as shown in SEQ ID NO:1.

Preferred nucleic acid fragments of the present invention include nucleic acid  
35 molecules encoding: a polypeptide comprising the TR10 receptor extracellular domain (amino acid residues from about 56 to about 212 in FIG. 1 or from about 1 to about

157 in SEQ ID NO:2); a polypeptide comprising the TR10 transmembrane domain (amino acid residues from about 213 to about 230 in FIG. 1 or from about 158 to about 175 in SEQ ID NO:2); a polypeptide comprising the TR10 intracellular domain (amino acid residues from about 231 to about 386 in FIG. 1 or from about 176 to about 331 in SEQ ID NO:2); and a polypeptide comprising the incomplete TR10 death domain (amino acid residues from about 353 to about 363 in FIG. 1 or from about 298 to about 308 in SEQ ID NO:2). Since the location of these domains have been predicted by computer analysis, one of ordinary skill would appreciate that the amino acid residues constituting these domains may vary slightly (e.g., by about 1 to 15 amino acid residues) depending on the criteria used to define each domain.

Preferred nucleic acid fragments of the invention encode a full-length TR10 polypeptide lacking the nucleotides encoding the amino terminal methionine (nucleotides 109-111 in SEQ ID NO:1), as it is known that the methionine is cleaved naturally and such sequences may be useful in genetically engineering TR10 expression vectors. Polypeptides encoded by such polynucleotides are also contemplated by the invention.

Preferred nucleic acid fragments of the present invention further include nucleic acid molecules encoding epitope-bearing portions of the TR10 receptor protein. In particular, such nucleic acid fragments of the present invention include nucleic acid molecules encoding: a polypeptide comprising amino acid residues from about 57 to about 113 in Figure 1 (corresponding to about amino acid 2 to about 58 in SEQ ID NO:2); a polypeptide comprising amino acid residues from about 130 to about 197 in Figure 1 (corresponding to about amino acid 75 to about 142 in SEQ ID NO:2); and a polypeptide comprising amino acid residues from about 250 to about 283 in Figure 1 (corresponding to about amino acid 195 to about 228 in SEQ ID NO:2). The inventors have determined that the above polypeptide fragments are antigenic regions of the TR10 protein. Methods for determining other such epitope-bearing portions of the TR10 protein are described in detail below.

In another aspect, the invention provides an isolated nucleic acid molecule comprising a polynucleotide which hybridizes under stringent hybridization conditions to a portion of the polynucleotide in a nucleic acid molecule of the invention described above, for instance, the cDNA clones contained in ATCC Deposit No. 209040. By "stringent hybridization conditions" is intended overnight incubation at 42°C in a solution comprising: 50% formamide, 5x SSC (150 mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10% dextran



sulfate, and 20 g/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1x SSC at about 65°C.

By a polynucleotide which hybridizes to a "portion" of a polynucleotide is intended a polynucleotide (either DNA or RNA) hybridizing to at least about 15 nucleotides (nt), and more preferably at least about 20 nt, still more preferably at least about 30 nt, and even more preferably about 30-70 nt of the reference polynucleotide. These are useful as diagnostic probes and primers as discussed above and in more detail below.

By a portion of a polynucleotide of "at least 20 nt in length," for example, is intended 20 or more contiguous nucleotides from the nucleotide sequence of the reference polynucleotide (e.g., the deposited cDNA or the nucleotide sequence as shown in SEQ ID NO:1).

Of course, a polynucleotide which hybridizes only to a poly A sequence (such as the 3' terminal poly(A) tract of the TR10 cDNA shown in SEQ ID NO:1), or to a complementary stretch of T (or U) residues, would not be included in a polynucleotide of the invention used to hybridize to a portion of a nucleic acid of the invention, since such a polynucleotide would hybridize to any nucleic acid molecule containing a poly (A) stretch or the complement thereof (e.g., practically any double-stranded cDNA clone).

As indicated, nucleic acid molecules of the present invention which encode a TR10 polypeptide may include, but are not limited to the coding sequence for the mature polypeptide, by itself; the coding sequence for the mature polypeptide and additional sequences, such as those encoding a leader or secretory sequence, such as a pre-, or pro- or prepro- protein sequence; the coding sequence of the mature polypeptide, with or without the aforementioned additional coding sequences, together with additional, non-coding sequences, including for example, but not limited to introns and non-coding 5' and 3' sequences, such as the transcribed, non-translated sequences that play a role in transcription, mRNA processing - including splicing and polyadenylation signals, for example - ribosome binding and stability of mRNA; additional coding sequence which codes for additional amino acids, such as those which provide additional functionalities. Thus, for instance, the polypeptide may be fused to a marker sequence, such as a peptide, which facilitates purification of the fused polypeptide. In certain preferred embodiments of this aspect of the invention, the marker sequence is a hexa-histidine peptide, such as the tag provided in a pQE vector (Qiagen, Inc.), among others, many of which are commercially available. As described in Gentz *et al.*, *Proc. Natl. Acad. Sci. USA* 86: 821-824 (1989), for instance, hexa-histidine provides for convenient purification of the fusion protein. The "HA" tag is



another peptide useful for purification which corresponds to an epitope derived from the influenza hemagglutinin protein, which has been described by Wilson *et al.*, *Cell* 37:767-778 (1984). As discussed below, other such fusion proteins include the TR10 receptor fused to Fc at the N- or C-terminus.

5       The present invention further relates to variants of the nucleic acid molecules of the present invention, which encode portions, analogs, or derivatives of the TR10 receptor. Variants may occur naturally, such as a natural allelic variant. By an "allelic variant" is intended one of several alternate forms of a gene occupying a given locus on a chromosome of an organism. *Genes II*, Lewin, B., ed., John Wiley & Sons, New  
10   York (1985). Non-naturally occurring variants may be produced using art-known mutagenesis techniques.

Such variants include those produced by nucleotide substitutions, deletions or additions which may involve one or more nucleotides. The variants may be altered in coding or non-coding regions or both. Alterations in the coding regions may produce  
15   conservative or non-conservative amino acid substitutions, deletions, or additions. Especially preferred among these are silent substitutions, additions, and deletions, which do not alter the properties and activities of the TR10 receptor or portions thereof. Also especially preferred in this regard are conservative substitutions.

Further embodiments of the invention include isolated nucleic acid molecules  
20   comprising a polynucleotide having a nucleotide sequence at least 90% identical, and more preferably at least 95%, 96%, 97%, 98%, or 99% identical to: (a) a nucleotide sequence encoding the polypeptide having the amino acid sequence in SEQ ID NO:2; (b) a nucleotide sequence encoding the polypeptide having the amino acid sequence in SEQ ID NO: 2, but lacking the amino terminal methionine; (c) a nucleotide sequence  
25   encoding the polypeptide having the amino acid sequence at positions about 1 to about 331 in SEQ ID NO:2; (d) a nucleotide sequence encoding the polypeptide having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. 209040; (e) a nucleotide sequence encoding the mature TR10 polypeptide having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No.  
30   209040; (f) a nucleotide sequence encoding the TR10 receptor extracellular domain; (g) a nucleotide sequence encoding the TR10 receptor transmembrane domain; (h) a nucleotide sequence encoding the TR10 receptor intracellular domain; (i) a nucleotide sequence encoding the TR10 receptor extracellular and intracellular domains with all or part of the transmembrane domain deleted; (j) a nucleotide sequence encoding the TR10  
35   receptor partial death domain; and (k) a nucleotide sequence complementary to any of the nucleotide sequences in (a), (b), (c), (d), (e), (f), (g), (h), (i), or (j) above.

By a polynucleotide having a nucleotide sequence at least, for example, 95% "identical" to a reference nucleotide sequence encoding a TR10 polypeptide is intended that the nucleotide sequence of the polynucleotide is identical to the reference sequence except that the polynucleotide sequence may include up to five point mutations per each 100 nucleotides of the reference nucleotide sequence encoding the TR10 polypeptide. In other words, to obtain a polynucleotide having a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. These mutations of the reference sequence may occur at the 5' or 3' terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence.

As a practical matter, whether any particular nucleic acid molecule is at least 90%, 95%, 96%, 97%, 98% or 99% identical to, for instance, the nucleotide sequence shown in SEQ ID NO:1 or to the nucleotide sequence of the deposited cDNA clone can be determined conventionally using known computer programs such as the Bestfit program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group, University Research Park, 575 Science Drive, Madison, WI 53711). Bestfit uses the local homology algorithm of Smith and Waterman, *Advances in Applied Mathematics* 2: 482-489 (1981), to find the best segment of homology between two sequences. When using Bestfit or any other sequence alignment program to determine whether a particular sequence is, for instance, 95% identical to a reference sequence according to the present invention, the parameters are set, of course, such that the percentage of identity is calculated over the full length of the reference nucleotide sequence and that gaps in homology of up to 5% of the total number of nucleotides in the reference sequence are allowed.

The present application is directed to nucleic acid molecules at least 90%, 95%, 96%, 97%, 98%, or 99% identical to the nucleic acid sequence shown in SEQ ID NO:1, or to the nucleic acid sequence of the deposited cDNA, irrespective of whether they encode a polypeptide having TR10 receptor activity. This is because even where a particular nucleic acid molecule does not encode a polypeptide having TR10 activity, one of skill in the art would still know how to use the nucleic acid molecule, for instance, as a hybridization probe or a polymerase chain reaction (PCR) primer. Uses of the nucleic acid molecules of the present invention that do not encode a polypeptide

having TR10 receptor activity include, *inter alia*: (1) isolating the TR10 receptor gene or allelic variants thereof in a cDNA library; (2) *in situ* hybridization (e.g., "FISH") to metaphase chromosomal spreads to provide precise chromosomal location of the TR10 receptor gene, as described in Verma *et al.*, *Human Chromosomes: A Manual of Basic Techniques*, Pergamon Press, New York (1988); and (3) Northern Blot analysis for detecting TR10 receptor mRNA expression in specific tissues.

Preferred, however, are nucleic acid molecules having sequences at least 90%, 95%, 96%, 97%, 98% or 99% identical to the nucleic acid sequence shown in SEQ ID NO:1, or to the nucleic acid sequence of the deposited cDNA which do, in fact, encode a polypeptide having TR10 receptor activity. By "a polypeptide having TR10 receptor activity" is intended polypeptides exhibiting activity similar, but not necessarily identical, to an activity of the TR10 receptor of the invention (either the full-length protein or, preferably, the mature protein), as measured in a particular biological assay. For example, TR10 receptor activity can be measured using the cell death assays performed essentially as previously described (A.M. Chinnaiyan *et al.*, *Cell* 81: 505-512 (1995); M.P. Boldin *et al.*, *J. Biol. Chem.* 270:7795-8(1995); F.C. Kischkel *et al.*, *EMBO* 14:5579-5588 (1995); A.M. Chinnaiyan *et al.*, *J. Biol. Chem.* 271: 4961-4965 (1996)) and as set forth in Example 5, below. In MCF7 cells, plasmids encoding full-length TR10 or a candidate death domain containing receptor are co-transfected with the pLantern reporter construct encoding green fluorescent protein. Nuclei of cells transfected with TR10 will exhibit apoptotic morphology as assessed by DAPI staining. Similar to TNFR-1 and Fas/APO-1 (M. Muzio *et al.*, *Cell* 85:817-827 (1996); M. P. Boldin *et al.*, *Cell* 85:803-815 (1996); M. Tewari *et al.*, *J. Biol. Chem.* 270:3255-60 (1995)), TR10-induced apoptosis is blocked by the inhibitors of ICE-like proteases, CrmA and z-VAD-fmk. In addition, apoptosis induced by TR10 is also blocked by dominant negative versions of FADD (FADD-DN) or FLICE (FLICE-DN/MACHa1C360S).

Of course, due to the degeneracy of the genetic code, one of ordinary skill in the art will immediately recognize that a large number of the nucleic acid molecules having a sequence at least 90%, 95%, 96%, 97%, 98%, or 99% identical to the nucleic acid sequence of the deposited cDNA or the nucleic acid sequence shown in SEQ ID NO:1 will encode a polypeptide "having TR10 receptor activity." In fact, since degenerate variants of these nucleotide sequences all encode the same polypeptide, this will be clear to the skilled artisan even without performing the above described comparison assay. It will be further recognized in the art that, for such nucleic acid molecules that are not degenerate variants, a reasonable number will also encode a polypeptide having TR10

receptor activity. This is because the skilled artisan is fully aware of amino acid substitutions that are either less likely or not likely to significantly effect protein function (e.g., replacing one aliphatic amino acid with a second aliphatic amino acid).

For example, guidance concerning how to make phenotypically silent amino acid substitutions is provided in J.U. Bowie *et al.*, "Deciphering the Message in Protein Sequences: Tolerance to Amino Acid Substitutions," *Science* 247:1306-1310 (1990), wherein the authors indicate that proteins are surprisingly tolerant of amino acid substitutions.

#### 10 ***Polynucleotide assays***

This invention is also related to the use of TR10 polynucleotides to detect complementary polynucleotides such as, for example, as a diagnostic reagent. Detection of a mutated form of TR10 associated with a dysfunction will provide a diagnostic tool that can add or define a diagnosis of a disease or susceptibility to a disease which results from under-expression over-expression or altered expression of TR10 or a soluble form thereof, such as, for example, tumors or autoimmune disease.

Individuals carrying mutations in the TR10 gene may be detected at the DNA level by a variety of techniques. Nucleic acids for diagnosis may be obtained from a patient's cells, such as from blood, urine, saliva, tissue biopsy and autopsy material. The genomic DNA may be used directly for detection or may be amplified enzymatically by using PCR prior to analysis. (Saiki *et al.*, *Nature* 324:163-166 (1986)). RNA or cDNA may also be used in the same ways. As an example, PCR primers complementary to the nucleic acid encoding TR10 can be used to identify and analyze TR10 expression and mutations. For example, deletions and insertions can be detected by a change in size of the amplified product in comparison to the normal genotype. Point mutations can be identified by hybridizing amplified DNA to radiolabeled TR10 RNA or alternatively, radiolabeled TR10 antisense DNA sequences. Perfectly matched sequences can be distinguished from mismatched duplexes by RNase A digestion or by differences in melting temperatures.

Sequence differences between a reference gene and genes having mutations also may be revealed by direct DNA sequencing. In addition, cloned DNA segments may be employed as probes to detect specific DNA segments. The sensitivity of such methods can be greatly enhanced by appropriate use of PCR or another amplification method. For example, a sequencing primer is used with double-stranded PCR product or a single-stranded template molecule generated by a modified PCR. The sequence

determination is performed by conventional procedures with radiolabeled nucleotide or by automatic sequencing procedures with fluorescent-tags.

Genetic testing based on DNA sequence differences may be achieved by detection of alteration in electrophoretic mobility of DNA fragments in gels, with or without denaturing agents. Small sequence deletions and insertions can be visualized by high resolution gel electrophoresis. DNA fragments of different sequences may be distinguished on denaturing formamide gradient gels in which the mobilities of different DNA fragments are retarded in the gel at different positions according to their specific melting or partial melting temperatures (see, e.g., Myers *et al.*, *Science* 230:1242 (1985)).

Sequence changes at specific locations also may be revealed by nuclease protection assays, such as RNase and S1 protection or the chemical cleavage method (e.g., Cotton *et al.*, *Proc. Natl. Acad. Sci. USA* 85: 4397-4401 (1985)).

Thus, the detection of a specific DNA sequence may be achieved by methods such as hybridization, RNase protection, chemical cleavage, direct DNA sequencing or the use of restriction enzymes, (e.g., restriction fragment length polymorphisms ("RFLP") and Southern blotting of genomic DNA.

In addition to more conventional gel-electrophoresis and DNA sequencing, mutations also can be detected by *in situ* analysis.

### **Vectors and Host Cells**

The present invention also relates to vectors which include the isolated DNA molecules of the present invention, host cells which are genetically engineered with the recombinant vectors of the invention and the production of TR10 polypeptides or fragments thereof by recombinant techniques.

The polynucleotides may be joined to a vector containing a selectable marker for propagation in a host. Generally, a plasmid vector is introduced in a precipitate, such as a calcium phosphate precipitate, or in a complex with a charged lipid. If the vector is a virus, it may be packaged *in vitro* using an appropriate packaging cell line and then transduced into host cells.

The DNA insert should be operatively linked to an appropriate promoter, such as the phage lambda PL promoter, the *E. coli lac*, *trp* and *tac* promoters, the SV40 early and late promoters and promoters of retroviral LTRs, to name a few. Other suitable promoters will be known to the skilled artisan. The expression constructs will further contain sites for transcription initiation, termination and, in the transcribed region, a ribosome binding site for translation. The coding portion of the mature transcripts



expressed by the constructs will preferably include a translation initiating at the beginning and a termination codon (UAA, UGA or UAG) appropriately positioned at the end of the polypeptide to be translated.

As indicated, the expression vectors will preferably include at least one  
5 selectable marker. Such markers include dihydrofolate reductase or neomycin resistance for eukaryotic cell culture and tetracycline or ampicillin resistance genes for culturing in *E. coli* and other bacteria. Representative examples of appropriate hosts include, but are not limited to, bacterial cells, such as *E. coli*, *Streptomyces* and *Salmonella typhimurium* cells; fungal cells, such as yeast cells; insect cells such as  
10 *Drosophila* S2 and *Spodoptera* Sf9 cells; animal cells such as CHO, COS and Bowes melanoma cells; and plant cells. Appropriate culture mediums and conditions for the above-described host cells are known in the art.

Among vectors preferred for use in bacteria include pQE70, pQE60 and pQE-9, available from Qiagen; pBS vectors, Phagescript vectors, Bluescript vectors, pNH8A,  
15 pNH16a, pNH18A, pNH46A, available from Stratagene; and ptrc99a, pKK223-3, pKK233-3, pDR540, pRIT5 available from Pharmacia. Among preferred eukaryotic vectors are pWLNEO, pSV2CAT, pOG44, pXT1 and pSG available from Stratagene; and pSVK3, pBPV, pMSG and pSVL available from Pharmacia. Other suitable vectors will be readily apparent to the skilled artisan.

20 Introduction of the construct into the host cell can be effected by calcium phosphate transfection, DEAE-dextran mediated transfection, cationic lipid-mediated transfection, electroporation, transduction, infection or other methods. Such methods are described in many standard laboratory manuals, such as Davis *et al.*, *Basic Methods In Molecular Biology* (1986).

25 The polypeptide may be expressed in a modified form, such as a fusion protein, and may include not only secretion signals but also additional heterologous functional regions. Thus, for instance, a region of additional amino acids, particularly charged amino acids, may be added to the N-terminus of the polypeptide to improve stability and persistence in the host cell, during purification or during subsequent handling and  
30 storage. Also, peptide moieties may be added to the polypeptide to facilitate purification. Such regions may be removed prior to final preparation of the polypeptide. The addition of peptide moieties to polypeptides to engender secretion or excretion, to improve stability and to facilitate purification, among others, are familiar and routine techniques in the art. A preferred fusion protein comprises a heterologous  
35 region from immunoglobulin that is useful to solubilize proteins. For example, EP-A-O 464 533 (Canadian counterpart 2045869) discloses fusion proteins comprising various



portions of constant region of immunoglobulin molecules together with another human protein or part thereof. In many cases, the Fc part in a fusion protein is thoroughly advantageous for use in therapy and diagnosis and thus results, for example, in improved pharmacokinetic properties (EP-A 0232 262). On the other hand, for some uses, it would be desirable to be able to delete the Fc part after the fusion protein has been expressed, detected and purified in the advantageous manner described. This is the case when the Fc portion proves to be a hindrance to use in therapy and diagnosis, for example, when the fusion protein is to be used as an antigen for immunizations. In drug discovery, for example, human proteins, such as the hIL5-receptor, have been fused with Fc portions for the purpose of high-throughput screening assays to identify antagonists of hIL-5. See, D. Bennett *et al.*, *Journal of Molecular Recognition* 8:52-58 (1995) and K. Johanson *et al.*, *The Journal of Biological Chemistry* 270:16:9459-9471 (1995).

The TR10 receptor can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography ("HPLC") is employed for purification.

Polypeptides of the present invention include naturally purified products, products of chemical synthetic procedures, and products produced by recombinant techniques from a prokaryotic or eukaryotic host, including, for example, bacterial, yeast, higher plant, insect and mammalian cells. Depending upon the host employed in a recombinant production procedure, the polypeptides of the present invention may be glycosylated or non-glycosylated. In addition, polypeptides of the invention may also include an initial modified methionine residue, in some cases as a result of host-mediated processes.

TR10 receptor polynucleotides and polypeptides may be used in accordance with the present invention for a variety of applications, particularly those that make use of the chemical and biological properties of TR10. Among these are applications in treatment of tumors, resistance to parasites, bacteria and viruses, to induce proliferation of T-cells, endothelial cells and certain hematopoietic cells, to treat restenosis, graft vs. host disease, to regulate anti-viral responses and to prevent certain autoimmune diseases after stimulation of TR10 by an agonist. Additional applications relate to diagnosis and to treatment of disorders of cells, tissues and organisms. These aspects of the invention are discussed further below.

### ***TR10 Receptor Polypeptides and Fragments***

The invention further provides an isolated TR10 polypeptide having the amino acid sequence encoded by the deposited cDNA, or the amino acid sequence in SEQ ID NO:2, or a peptide or polypeptide comprising a portion of the above polypeptides.

It will be recognized in the art that some amino acid sequences of TR10 can be varied without significant effect on the structure or function of the protein. If such differences in sequence are contemplated, it should be remembered that there will be critical areas on the protein which determine activity. Thus, the invention further includes variations of the TR10 receptor, which show substantial TR10 receptor activity or which include regions of TR10 proteins, such as the protein portions discussed below. Such mutants include deletions, insertions, inversions, repeats, and type substitutions. As indicated above, guidance concerning which amino acid changes are likely to be phenotypically silent can be found in J.U. Bowie *et al.*, *Science* 247:1306-1310 (1990).

Thus, the fragment, derivative, or analog of the polypeptide of SEQ ID NO:2, or that encoded by the deposited cDNA, may be (i) one in which at least one or more of the amino acid residues are substituted with a conserved or non-conserved amino acid residue (preferably a conserved amino acid residue(s), and more preferably at least one but less than ten conserved amino acid residues) and such substituted amino acid residue may or may not be one encoded by the genetic code, or (ii) one in which one or more of the amino acid residues includes a substituent group, or (iii) one in which the mature polypeptide is fused with another compound, such as a compound to increase the half-life of the polypeptide (for example, polyethylene glycol), or (iv) one in which the additional amino acids are fused to the mature polypeptide, such as an IgG Fc fusion region peptide or leader or secretory sequence or a sequence which is employed for purification of the mature polypeptide or a proprotein sequence. Such fragments, derivatives and analogs are deemed to be within the scope of those skilled in the art from the teachings herein.

Of particular interest are substitutions of charged amino acids with another charged amino acid and with neutral or negatively charged amino acids. The latter results in proteins with reduced positive charge to improve the characteristics of the TR10 receptor protein. The prevention of aggregation is highly desirable. Aggregation of proteins not only results in a loss of activity but can also be problematic when preparing pharmaceutical formulations, because they can be immunogenic. (Pinckard *et al.*, *Clin Exp. Immunol.* 2:331-340 (1967); Robbins *et al.*, *Diabetes* 36:838-845

(1987); Cleland *et al.* *Crit. Rev. Therapeutic Drug Carrier Systems* 10:307-377 (1993)).

The replacement of amino acids can also change the selectivity of binding to cell surface receptors. Ostade *et al.*, *Nature* 361:266-268 (1993), describes certain mutations resulting in selective binding of TNF- to only one of the two known types of TNF receptors. Thus, the TR10 receptor of the present invention may include one or more amino acid substitutions, deletions, or additions, either from natural mutations or human manipulation.

As indicated, changes are preferably of a minor nature, such as conservative amino acid substitutions that do not significantly affect the folding or activity of the protein (see Table 1).

**TABLE 1. Conservative Amino Acid Substitutions**

Aromatic	Phenylalanine Tryptophan Tyrosine
Hydrophobic	Leucine Isoleucine Valine
Polar	Glutamine Asparagine
Basic	Arginine Lysine Histidine
Acidic	Aspartic Acid Glutamic Acid
Small	Alanine Serine Threonine Methionine Glycine

15

Amino acids in the TR10 protein of the present invention that are essential for function can be identified by methods known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham and Wells, *Science* 244:1081-1085 (1989)). The latter procedure introduces single alanine mutations at every residue in the molecule. The resulting mutant molecules are then tested for biological activity such as receptor binding or *in vitro* proliferative activity. Sites that

20

are critical for ligand-receptor binding can also be determined by structural analysis such as crystallization, nuclear magnetic resonance or photoaffinity labeling (Smith *et al.*, *J. Mol. Biol.* 224:899-904 (1992) and de Vos *et al.* *Science* 255:306-312 (1992)).

The polypeptides of the present invention are preferably provided in an isolated form. By "isolated polypeptide" is intended a polypeptide removed from its native environment. Thus, a polypeptide produced and/or contained within a recombinant host cell is considered isolated for purposes of the present invention. Also intended as an "isolated polypeptide" are polypeptides that have been purified, partially or substantially, from a recombinant host cell. For example, a recombinantly produced version of the TR10 polypeptide can be substantially purified by the one-step method described in Smith and Johnson, *Gene* 67:31-40 (1988).

The polypeptides of the present invention include the polypeptide encoded by the deposited cDNA including the leader; the mature polypeptide encoded by the deposited cDNA minus the leader (i.e., the mature protein); a polypeptide comprising amino acids about - 55 to about 331 in SEQ ID NO:2; a polypeptide comprising amino acids about - 54 to about 331 in SEQ ID NO:2; a polypeptide comprising amino acids about 1 to about 331 in SEQ ID NO:2; a polypeptide comprising the extracellular domain; a polypeptide comprising the transmembrane domain; a polypeptide comprising the intracellular domain; a polypeptide comprising the extracellular and intracellular domains with all or part of the transmembrane domain deleted; and a polypeptide comprising the partial death domain; as well as polypeptides which are at least 80% identical, more preferably at least 90% or 95% identical, still more preferably at least 96%, 97%, 98%, or 99% identical to the polypeptides described above, and also include portions of such polypeptides with at least 30 amino acids and more preferably at least 50 amino acids.

By a polypeptide having an amino acid sequence at least, for example, 95% "identical" to a reference amino acid sequence of a TR10 polypeptide is intended that the amino acid sequence of the polypeptide is identical to the reference sequence except that the polypeptide sequence may include up to five amino acid alterations per each 100 amino acids of the reference amino acid of the TR10 receptor. In other words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a reference amino acid sequence, up to 5% of the amino acid residues in the reference sequence may be deleted or substituted with another amino acid, or a number of amino acids up to 5% of the total amino acid residues in the reference sequence may be inserted into the reference sequence. These alterations of the reference sequence may occur at the amino or carboxy terminal positions of the reference amino acid sequence or anywhere between

those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

As a practical matter, whether any particular polypeptide is at least 90%, 95%, 96%, 97%, 98%, or 99% identical to, for instance, the amino acid sequence shown in SEQ ID NO:2, or to the amino acid sequence encoded by the deposited cDNA clone, can be determined conventionally using known computer programs such the Bestfit program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group, University Research Park, 575 Science Drive, Madison, WI 53711). When using Bestfit or any other sequence alignment program to determine whether a particular sequence is, for instance, 95% identical to a reference sequence according to the present invention, the parameters are set, of course, such that the percentage of identity is calculated over the full length of the reference amino acid sequence and that gaps in homology of up to 5% of the total number of amino acid residues in the reference sequence are allowed.

Particularly, N-terminal deletions of the TR10 polypeptide can be described by the general formula m-157, where m is a number from 1-156, corresponding to the position of amino acid identified in SEQ ID NO:2. Preferably, N-terminal deletions of the TR10 polypeptide of the invention shown as SEQ ID NO:2 include polypeptides comprising the amino acid sequence of residues: T-2 to Y-157; I-3 to Y-157; P-4 to Y-157; R-5 to Y-157; Q-6 to Y-157; D-7 to Y-157; E-8 to Y-157; V-9 to Y-157; P-10 to Y-157; Q-11 to Y-157; Q-12 to Y-157; T-13 to Y-157; V-14 to Y-157; A-15 to Y-157; P-16 to Y-157; Q-17 to Y-157; Q-18 to Y-157; Q-19 to Y-157; R-20 to Y-157; R-21 to Y-157; S-22 to Y-157; L-23 to Y-157; K-24 to Y-157; E-25 to Y-157; E-26 to Y-157; E-27 to Y-157; C-28 to Y-157; P-29 to Y-157; A-30 to Y-157; G-31 to Y-157; S-32 to Y-157; H-33 to Y-157; R-34 to Y-157; S-35 to Y-157; E-36 to Y-157; Y-37 to Y-157; T-38 to Y-157; G-39 to Y-157; A-40 to Y-157; C-41 to Y-157; N-42 to Y-157; P-43 to Y-157; C-44 to Y-157; T-45 to Y-157; E-46 to Y-157; G-47 to Y-157; V-48 to Y-157; D-49 to Y-157; Y-50 to Y-157; T-51 to Y-157; I-52 to Y-157; A-53 to Y-157; S-54 to Y-157; N-55 to Y-157; N-56 to Y-157; L-57 to Y-157; P-58 to Y-157; S-59 to Y-157; C-60 to Y-157; L-61 to Y-157; L-62 to Y-157; C-63 to Y-157; T-64 to Y-157; V-65 to Y-157; C-66 to Y-157; K-67 to Y-157; S-68 to Y-157; G-69 to Y-157; Q-70 to Y-157; T-71 to Y-157; N-72 to Y-157; K-73 to Y-157; S-74 to Y-157; S-75 to Y-157; C-76 to Y-157; T-77 to Y-157; T-78 to Y-157; T-79 to Y-157; R-80 to Y-157; D-81 to Y-157; T-82 to Y-157; V-83 to Y-157; C-84 to Y-157; Q-85 to Y-157; C-86 to Y-157; E-87 to Y-157; K-88 to Y-157; G-89 to Y-157; S-90 to Y-157; F-91 to Y-157; Q-92 to Y-157; D-93 to Y-157; K-94 to Y-157; N-95 to Y-157; S-96 to Y-157; P-97 to Y-157; E-98 to Y-157; M-99 to Y-157; C-100 to Y-157; R-101 to Y-157; T-102 to Y-157; C-103 to Y-157; R-104 to Y-157; T-105 to Y-157; G-106 to Y-157; C-107 to Y-157; P-108 to Y-157; R-109 to Y-157; G-110 to Y-157; M-111 to Y-157; V-112 to Y-157; K-

113 to Y-157; V-114 to Y-157; S-115 to Y-157; N-116 to Y-157; C-117 to Y-157; T-118 to Y-157; P-119 to Y-157; R-120 to Y-157; S-121 to Y-157; D-122 to Y-157; I-123 to Y-157; K-124 to Y-157; C-125 to Y-157; K-126 to Y-157; N-127 to Y-157; E-128 to Y-157; S-129 to Y-157; A-130 to Y-157; A-131 to Y-157; S-132 to Y-157; S-133 to Y-157; T-134 to Y-157; G-135 to Y-157; K-136 to Y-157; T-137 to Y-157; P-138 to Y-157; A-139 to Y-157; A-140 to Y-157; E-141 to Y-157; E-142 to Y-157; T-143 to Y-157; V-144 to Y-157; T-145 to Y-157; T-146 to Y-157; I-147 to Y-157; L-148 to Y-157; G-149 to Y-157; M-150 to Y-157; L-151 to Y-157; A-152 to Y-157; of SEQ ID NO:2.

Moreover, C-terminal deletions of the TR10 polypeptide can also be described by the general formula 1-n, where n is a number from 2-156, corresponding to the position of amino acid identified in SEQ ID NO:2. Preferably, C-terminal deletions of the TR10 polypeptide of the invention shown as SEQ ID NO:2 include polypeptides comprising the amino acid sequence of residues: A-1 to H-156; A-1 to Y-155; A-1 to P-154; A-1 to S-153; A-1 to A-152; A-1 to L-151; A-1 to M-150; A-1 to G-149; A-1 to L-148; A-1 to I-147; A-1 to T-146; A-1 to T-145; A-1 to V-144; A-1 to T-143; A-1 to E-142; A-1 to E-141; A-1 to A-140; A-1 to A-139; A-1 to P-138; A-1 to T-137; A-1 to K-136; A-1 to G-135; A-1 to T-134; A-1 to S-133; A-1 to S-132; A-1 to A-131; A-1 to A-130; A-1 to S-129; A-1 to E-128; A-1 to N-127; A-1 to K-126; A-1 to C-125; A-1 to K-124; A-1 to I-123; A-1 to D-122; A-1 to S-121; A-1 to R-120; A-1 to P-119; A-1 to T-118; A-1 to C-117; A-1 to N-116; A-1 to S-115; A-1 to V-114; A-1 to K-113; A-1 to V-112; A-1 to M-111; A-1 to G-110; A-1 to R-109; A-1 to P-108; A-1 to C-107; A-1 to G-106; A-1 to T-105; A-1 to R-104; A-1 to C-103; A-1 to T-102; A-1 to R-101; A-1 to C-100; A-1 to M-99; A-1 to E-98; A-1 to P-97; A-1 to S-96; A-1 to N-95; A-1 to K-94; A-1 to D-93; A-1 to Q-92; A-1 to F-91; A-1 to S-90; A-1 to G-89; A-1 to K-88; A-1 to E-87; A-1 to C-86; A-1 to Q-85; A-1 to C-84; A-1 to V-83; A-1 to T-82; A-1 to D-81; A-1 to R-80; A-1 to T-79; A-1 to T-78; A-1 to T-77; A-1 to C-76; A-1 to S-75; A-1 to S-74; A-1 to K-73; A-1 to N-72; A-1 to T-71; A-1 to Q-70; A-1 to G-69; A-1 to S-68; A-1 to K-67; A-1 to C-66; A-1 to V-65; A-1 to T-64; A-1 to C-63; A-1 to L-62; A-1 to L-61; A-1 to C-60; A-1 to S-59; A-1 to P-58; A-1 to L-57; A-1 to N-56; A-1 to N-55; A-1 to S-54; A-1 to A-53; A-1 to I-52; A-1 to T-51; A-1 to Y-50; A-1 to D-49; A-1 to V-48; A-1 to G-47; A-1 to E-46; A-1 to T-45; A-1 to C-44; A-1 to P-43; A-1 to N-42; A-1 to C-41; A-1 to A-40; A-1 to G-39; A-1 to T-38; A-1 to Y-37; A-1 to E-36; A-1 to S-35; A-1 to R-34; A-1 to H-33; A-1 to S-32; A-1 to G-31; A-1 to A-30; A-1 to P-29; A-1 to C-28; A-1 to E-27; A-1 to E-26; A-1 to E-25; A-1 to K-24; A-1 to L-23; A-1 to S-22; A-1 to R-21; A-1 to R-20; A-1 to Q-19; A-1 to Q-18; A-1 to Q-17; A-1 to P-16; A-1 to A-15; A-1 to V-14; A-1 to T-13; A-1 to Q-12; A-1 to Q-11; A-1 to P-10; A-1 to V-9; A-1 to E-8; A-1 to D-7; of SEQ ID NO:2.

For example, any of the above listed N- or C-terminal deletions can be combined to



produce a N- and C-terminal deleted TR10 polypeptide.

The invention also provides polypeptides having one or more amino acids deleted from both the amino and the carboxyl termini, which may be described generally as having residues n-m of SEQ ID NO:2, where n and m are integers as  
5 described above.

The polypeptide of the present invention could be used as a molecular weight marker on SDS-PAGE gels or on molecular sieve gel filtration columns using methods well known to those of skill in the art.

In another aspect, the invention provides a peptide or polypeptide comprising an  
10 epitope-bearing portion of a polypeptide of the invention. The epitope of this polypeptide portion is an immunogenic or antigenic epitope of a polypeptide described herein. An "immunogenic epitope" is defined as a part of a protein that elicits an antibody response when the whole protein is the immunogen. On the other hand, a region of a protein molecule to which an antibody can bind is defined as an "antigenic  
15 epitope." The number of immunogenic epitopes of a protein generally is less than the number of antigenic epitopes. See, for instance, Geysen *et al.*, *Proc. Natl. Acad. Sci. USA* 81:3998- 4002 (1983).

As to the selection of peptides or polypeptides bearing an antigenic epitope (i.e., that contain a region of a protein molecule to which an antibody can bind), it is well  
20 known in that art that relatively short synthetic peptides that mimic part of a protein sequence are routinely capable of eliciting an antiserum that reacts with the partially mimicked protein. See, for instance, J.G. Sutcliffe *et al.*, "Antibodies That React With Predetermined Sites on Proteins," *Science* 219:660-666 (1983). Peptides capable of eliciting protein-reactive sera are frequently represented in the primary sequence of a  
25 protein, can be characterized by a set of simple chemical rules, and are confined neither to immunodominant regions of intact proteins (i.e., immunogenic epitopes) nor to the amino or carboxyl terminals.

Antigenic epitope-bearing peptides and polypeptides of the invention are therefore useful to raise antibodies, including monoclonal antibodies, that bind  
30 specifically to a polypeptide of the invention. See, for instance, Wilson *et al.*, *Cell* 37:767-778 (1984) at 777. Antigenic epitope-bearing peptides and polypeptides of the invention preferably contain a sequence of at least seven, more preferably at least nine, and most preferably between at least about 15 to about 30 amino acids contained within the amino acid sequence of a polypeptide of the invention.

35 Non-limiting examples of antigenic polypeptides or peptides that can be used to generate TR10 receptor-specific antibodies include: a polypeptide comprising amino

acid residues from about 57 to about 113 in Figure 1 (2 to 58 in SEQ ID NO:2); a polypeptide comprising amino acid residues from about 130 to about 197 in Figure 1 (75 to 142 in SEQ ID NO:2); and a polypeptide comprising amino acid residues from about 250 to about 283 in Figure 1 (195 to 228 in SEQ ID NO:2). As indicated above, the inventors have determined that the above polypeptide fragments are antigenic regions of the TR10 receptor protein.

The epitope-bearing peptides and polypeptides of the invention may be produced by any conventional means. R.A. Houghten, "General Method for the Rapid Solid-phase Synthesis of Large Numbers of Peptides: Specificity of Antigen-Antibody Interaction at the Level of Individual Amino Acids," *Proc. Natl. Acad. Sci. USA* 82:5131-5135 (1985). This "Simultaneous Multiple Peptide Synthesis (SMPS)" process is further described in U.S. Patent No. 4,631,211 to Houghten *et al.* (1986).

As one of skill in the art will appreciate, TR10 receptor polypeptides of the present invention and the epitope-bearing fragments thereof, described above, can be combined with parts of the constant domain of immunoglobulins (IgG), resulting in chimeric polypeptides. These fusion proteins facilitate purification and show an increased half-life *in vivo*. This has been shown, e.g., for chimeric proteins consisting of the first two domains of the human CD4-polypeptide and various domains of the constant regions of the heavy or light chains of mammalian immunoglobulins (EPA 394,827; Traunecker *et al.*, *Nature* 331:84-86 (1988)). Fusion proteins that have a disulfide-linked dimeric structure due to the IgG part can also be more efficient in binding and neutralizing other molecules than the monomeric TR10 protein or protein fragment alone (Fountoulakis *et al.*, *J. Biochem.* 270:3958-3964 (1995)).

### ***Polypeptide assays***

The present invention also relates to diagnostic assays such as quantitative and diagnostic assays for detecting levels of TR10 receptor protein, or the soluble form thereof, in cells and tissues, including determination of normal and abnormal levels. Thus, for instance, a diagnostic assay in accordance with the invention for detecting over-expression of TR10, or soluble form thereof, compared to normal control tissue samples may be used to detect the presence of tumors, for example. Assay techniques that can be used to determine levels of a protein, such as a TR10 protein of the present invention, or a soluble form thereof, in a sample derived from a host are well-known to those of skill in the art. Such assay methods include radioimmunoassays, competitive-binding assays, Western Blot analysis and ELISA assays.

Assaying TR10 protein levels in a biological sample can occur using any art-known method. By "biological sample" is intended any biological sample obtained from an individual, cell line, tissue culture, or other source containing TR10 receptor protein or mRNA. Preferred for assaying TR10 protein levels in a biological sample are antibody-based techniques. For example, TR10 protein expression in tissues can be studied with classical immunohistological methods. (M. Jalkanen *et al.*, *J. Cell. Biol.* 101:976-985 (1985); M. Jalkanen *et al.*, *J. Cell. Biol.* 105:3087-3096 (1987)). Other antibody-based methods useful for detecting TR10 receptor gene expression include immunoassays, such as the enzyme linked immunosorbent assay (ELISA) and the radioimmunoassay (RIA).

Suitable labels are known in the art and include enzyme labels, such as glucose oxidase, radioisotopes, such as iodine ( $^{125}\text{I}$ ,  $^{121}\text{I}$ ), carbon ( $^{14}\text{C}$ ), sulphur ( $^{35}\text{S}$ ), tritium ( $^3\text{H}$ ), indium ( $^{112}\text{In}$ ), and technetium ( $^{99\text{m}}\text{Tc}$ ), and fluorescent labels, such as fluorescein and rhodamine, and biotin.

15

### ***Therapeutics***

The Tumor Necrosis Factor (TNF) family ligands are known to be among the most pleiotropic cytokines, inducing a large number of cellular responses, including cytotoxicity, anti-viral activity, immunoregulatory activities, and the transcriptional regulation of several genes (D.V. Goeddel *et al.*, "Tumor Necrosis Factors: Gene Structure and Biological Activities," *Symp. Quant. Biol.* 51:597-609 (1986), Cold Spring Harbor; B. Beutler and A. Cerami, *Annu. Rev. Biochem.* 57:505-518 (1988); L.J. Old, *Sci. Am.* 258:59-75 (1988); W. Fiers, *FEBS Lett.* 285:199-224 (1991)). The TNF-family ligands induce such various cellular responses by binding to TNF-family receptors, including the TR10 of the present invention. Cells which express the TR10 polypeptide and are believed to have a potent cellular response to TR10 ligands include fetal liver, PBL, lung, kidney, small intestine, colon, keratinocytes, endothelial cells, and monocyte activated tissue. By "a cellular response to a TNF-family ligand" is intended any genotypic, phenotypic, and/or morphologic change to a cell, cell line, tissue, tissue culture or patient that is induced by a TNF-family ligand. As indicated, such cellular responses include not only normal physiological responses to TNF-family ligands, but also diseases associated with increased apoptosis or the inhibition of apoptosis. Apoptosis-programmed cell death-is a physiological mechanism involved in the deletion of peripheral T lymphocytes of the immune system, and its dysregulation can lead to a number of different pathogenic processes (J.C. Ameisen, *AIDS* 8:1197-1213 (1994); P.H. Krammer *et al.*, *Curr. Opin. Immunol.* 6:279-289 (1994)).

35

Diseases associated with increased cell survival, or the inhibition of apoptosis, include cancers (such as follicular lymphomas, carcinomas with p53 mutations, and hormone-dependent tumors, such as breast cancer, prostate cancer, Kaposi's sarcoma and ovarian cancer); autoimmune disorders (such as systemic lupus erythematosus and immune-related glomerulonephritis rheumatoid arthritis); viral infections (such as herpes viruses, pox viruses and adenoviruses); inflammation; graft vs. host disease; acute graft rejection and chronic graft rejection. Diseases associated with increased apoptosis include AIDS; neurodegenerative disorders (such as Alzheimer's disease, Parkinson's disease, Amyotrophic lateral sclerosis, Retinitis pigmentosa, Cerebellar degeneration); myelodysplastic syndromes (such as aplastic anemia), ischemic injury (such as that caused by myocardial infarction, stroke and reperfusion injury), toxin-induced liver disease (such as that caused by alcohol), septic shock, cachexia and anorexia.

Thus, in one aspect, the present invention is directed to a method for enhancing apoptosis induced by a TNF-family ligand, which involves administering to a cell which expresses the TR10 polypeptide an effective amount of TR10 ligand, analog or an agonist capable of increasing TR10 mediated signaling. Preferably, TR10 mediated signaling is increased to treat a disease wherein decreased apoptosis or decreased cytokine and adhesion molecule expression is exhibited. An agonist can include soluble forms of TR10 and monoclonal antibodies directed against the TR10 polypeptide.

In a further aspect, the present invention is directed to a method for inhibiting apoptosis induced by a TNF-family ligand, which involves administering to a cell which expresses the TR10 polypeptide an effective amount of an antagonist capable of decreasing TR10 mediated signaling. Preferably, TR10 mediated signaling is decreased to treat a disease wherein increased apoptosis or NFkB expression is exhibited. An antagonist can include soluble forms of TR10 and monoclonal antibodies directed against the TR10 polypeptide.

By "agonist" is intended naturally occurring and synthetic compounds capable of enhancing or potentiating apoptosis. By "antagonist" is intended naturally occurring and synthetic compounds capable of inhibiting apoptosis. Whether any candidate "agonist" or "antagonist" of the present invention can enhance or inhibit apoptosis can be determined using art-known TNF-family ligand/receptor cellular response assays, including those described in more detail below.

One such screening procedure involves the use of melanophores which are transfected to express the receptor of the present invention. Such a screening technique is described in PCT WO 92/01810, published February 6, 1992. Such an assay may

be employed, for example, for screening for a compound which inhibits (or enhances) activation of the receptor polypeptide of the present invention by contacting the melanophore cells which encode the receptor with both a TNF-family ligand and the candidate antagonist (or agonist). Inhibition or enhancement of the signal generated by the ligand indicates that the compound is an antagonist or agonist of the ligand/receptor signaling pathway.

Other screening techniques include the use of cells which express the receptor (for example, transfected CHO cells) in a system which measures extracellular pH changes caused by receptor activation. For example, compounds may be contacted with a cell which expresses the receptor polypeptide of the present invention and a second messenger response, e.g., signal transduction or pH changes, may be measured to determine whether the potential compound activates or inhibits the receptor.

Another such screening technique involves introducing RNA encoding the receptor into *Xenopus* oocytes to transiently express the receptor. The receptor oocytes may then be contacted with the receptor ligand and a compound to be screened, followed by detection of inhibition or activation of a calcium signal in the case of screening for compounds which are thought to inhibit activation of the receptor.

Another screening technique well known in the art involves expressing in cells a construct wherein the receptor is linked to a phospholipase C or D. Exemplary cells include endothelial cells, smooth muscle cells, embryonic kidney cells, etc. The screening may be accomplished as hereinabove described by detecting activation of the receptor or inhibition of activation of the receptor from the phospholipase signal.

Another method involves screening for compounds which inhibit activation of the receptor polypeptide of the present invention antagonists by determining inhibition of binding of labeled ligand to cells which have the receptor on the surface thereof. Such a method involves transfecting a eukaryotic cell with DNA encoding the receptor such that the cell expresses the receptor on its surface and contacting the cell with a compound in the presence of a labeled form of a known ligand. The ligand can be labeled, e.g., by radioactivity. The amount of labeled ligand bound to the receptors is measured, e.g., by measuring radioactivity of the receptors. If the compound binds to the receptor as determined by a reduction of labeled ligand which binds to the receptors, the binding of labeled ligand to the receptor is inhibited.

Further screening assays for agonists and antagonists of the present invention are described in L.A. Tartaglia and D.V. Goeddel, *J. Biol. Chem.* 267:4304-4307(1992).



Thus, in a further aspect, a screening method is provided for determining whether a candidate agonist or antagonist is capable of enhancing or inhibiting a cellular response to a TNF-family ligand. The method involves contacting cells which express the TR10 polypeptide with a candidate compound and a TNF-family ligand, assaying a cellular response, and comparing the cellular response to a standard cellular response, the standard being assayed when contact is made with the ligand in absence of the candidate compound, whereby an increased cellular response over the standard indicates that the candidate compound is an agonist of the ligand/receptor signaling pathway and a decreased cellular response compared to the standard indicates that the candidate compound is an antagonist of the ligand/receptor signaling pathway. By "assaying a cellular response" is intended qualitatively or quantitatively measuring a cellular response to a candidate compound and/or a TNF-family ligand (e.g., determining or estimating an increase or decrease in T cell proliferation or tritiated thymidine labeling). By the invention, a cell expressing the TR10 polypeptide can be contacted with either an endogenous or exogenously administered TNF-family ligand.

Antagonist according to the present invention include naturally occurring and synthetic compounds such as, for example, TNF family ligand peptide fragments, transforming growth factor, neurotransmitters (such as glutamate, dopamine, *N*-methyl-D-aspartate), tumor suppressors (p53), cytolytic T cells and antimetabolites. Preferred agonists include chemotherapeutic drugs such as, for example, cisplatin, doxorubicin, bleomycin, cytosine arabinoside, nitrogen mustard, methotrexate and vincristine. Others include ethanol and  $\beta$ -amyloid peptide. (*Science* 267:1457-1458 (1995)). Further preferred agonists include polyclonal and monoclonal antibodies raised against the TR10 polypeptide, or a fragment thereof. Such agonist antibodies raised against a TNF-family receptor are disclosed in L.A. Tartaglia *et al.*, *Proc. Natl. Acad. Sci. USA* 88:9292-9296 (1991); and L.A. Tartaglia and D.V. Goeddel, *J. Biol. Chem.* 267:4304-4307 (1992). See, also, PCT Application WO 94/09137.

Agonists according to the present invention include naturally occurring and synthetic compounds such as, for example, the CD40 ligand, neutral amino acids, zinc, estrogen, androgens, viral genes (such as Adenovirus *E1B*, Baculovirus *p35* and *IAP*, Cowpox virus *crmA*, Epstein-Barr virus *BHRF1*, *LMP-1*, African swine fever virus *LMW5-HL*, and Herpesvirus  $\gamma$ 1 34.5), calpain inhibitors, cysteine protease inhibitors, and tumor promoters (such as PMA, Phenobarbital, and  $\gamma$ -Hexachlorocyclohexane).

Other potential antagonists include antisense molecules. Antisense technology can be used to control gene expression through antisense DNA or RNA or through triple-helix formation. Antisense techniques are discussed, for example, in Okano, *J.*



*Neurochem.* 56:560 (1991); *Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression*, CRC Press, Boca Raton, FL (1988). Triple helix formation is discussed in, for instance Lee *et al.*, *Nucleic Acids Research* 6:3073 (1979); Cooney *et al.*, *Science* 241:456 (1988); and Dervan *et al.*, *Science* 251:1360 (1991). The methods are  
5 based on binding of a polynucleotide to a complementary DNA or RNA.

For example, the 5' coding portion of a polynucleotide that encodes the mature polypeptide of the present invention may be used to design an antisense RNA oligonucleotide of from about 10 to 40 base pairs in length. A DNA oligonucleotide is designed to be complementary to a region of the gene involved in transcription thereby  
10 preventing transcription and the production of the receptor. The antisense RNA oligonucleotide hybridizes to the mRNA *in vivo* and blocks translation of the mRNA molecule into receptor polypeptide. The oligonucleotides described above can also be delivered to cells such that the antisense RNA or DNA may be expressed *in vivo* to inhibit production of the receptor.

15 An agonists according to the present invention include soluble forms of TR10, i.e., TR10 fragments that include the ligand binding domain from the extracellular region of the full length receptor. Such soluble forms of the receptor, which may be naturally occurring or synthetic, antagonize TR10 mediated signaling by competing with the cell surface TR10 for binding to TNF-family ligands. However, soluble TR10  
20 may bind to apoptosis inducing ligands such as TRAIL and more effectively compete for TRAIL binding reducing the available TRAIL for binding to receptors with functional death domains. Thus, soluble forms of the receptor that include the ligand binding domain are novel cytokines capable of inhibiting apoptosis induced by TNF-family ligands. These are preferably expressed as dimers or trimers, since these have  
25 been shown to be superior to monomeric forms of soluble receptor as antagonists, e.g., IgGFc-TNF receptor family fusions. Other such cytokines are known in the art and include Fas B (a soluble form of the mouse Fas receptor) that acts physiologically to limit apoptosis induced by Fas ligand (D.P. Hughes and I.N. Crispe, *J. Exp. Med.* 182:1395-1401 (1995)).

30 The term "antibody" (Ab) or "monoclonal antibody" (mAb) as used herein is meant to include intact molecules as well as fragments thereof (such as, e.g., Fab and F(ab')<sub>2</sub> fragments) which are capable of binding an antigen. Fab and F(ab')<sub>2</sub> fragments lack the Fc fragment of intact antibody, clear more rapidly from the circulation, and may have less non-specific tissue binding of an intact antibody (Wahl *et al.*, *J. Nucl. Med.* 24:316-325 (1983)).  
35

Antibodies according to the present invention may be prepared by any of a variety of methods using TR10 immunogens of the present invention. As indicated, such TR10 immunogens include the full length TR10 polypeptide (which may or may not include the leader sequence) and TR10 polypeptide fragments such as the ligand  
5 binding domain, the transmembrane domain, the intracellular domain and the incomplete death domain.

Proteins and other compounds which bind the TR10 domains are also candidate agonists and antagonists according to the present invention. Such binding compounds can be "captured" using the yeast two-hybrid system (Fields and Song, *Nature*  
10 340:245-246 (1989)). A modified version of the yeast two-hybrid system has been described by Roger Brent and his colleagues (J. Gyuris, *Cell* 75:791-803 (1993); A.S. Zervos *et al.*, *Cell* 72:223-232 (1993)). Preferably, the yeast two-hybrid system is used according to the present invention to capture compounds which bind to either the TR10 ligand binding domain or to the TR10 intracellular domain. Such compounds are  
15 good candidate agonists and antagonists of the present invention.

By a "TNF-family ligand" is intended naturally occurring, recombinant, and synthetic ligands that are capable of binding to a member of the TNF receptor family and inducing the ligand/receptor signaling pathway. Members of the TNF ligand family include, but are not limited to TR10 ligands including TRAIL, TNF- $\alpha$ , lymphotoxin- $\alpha$   
20 (LT- $\alpha$ , also known as TNF- $\beta$ ), LT- $\beta$  (found in complex heterotrimer LT- $\alpha$ 2- $\beta$ ), FasL, CD40, CD27, CD30, 4-1BB, OX40, and nerve growth factor (NGF).

Representative therapeutic applications of the present invention are discussed in more detail below. The state of immunodeficiency that defines AIDS is secondary to a decrease in the number and function of CD4<sup>+</sup> T-lymphocytes. Recent reports estimate  
25 the daily loss of CD4<sup>+</sup> T cells to be between  $3.5 \times 10^7$  and  $2 \times 10^9$  cells (X. Wei *et al.*, *Nature* 373:117-122 (1995)). One cause of CD4<sup>+</sup> T cell depletion in the setting of HIV infection is believed to be HIV-induced apoptosis. Indeed, HIV-induced apoptotic cell death has been demonstrated not only *in vitro* but also, more importantly, in infected individuals (J.C. Ameisen, *AIDS* 8:1197-1213 (1994); T.H. Finkel and N.K. Banda,  
30 *Curr. Opin. Immunol.* 6:605-615(1995); C.A. Muro-Cacho *et al.*, *J. Immunol.* 154:5555-5566 (1995)). Furthermore, apoptosis and CD4<sup>+</sup> T-lymphocyte depletion is tightly correlated in different animal models of AIDS (T. Brunner *et al.*, *Nature* 373:441-444 (1995); M.L. Gougeon *et al.*, *AIDS Res. Hum. Retroviruses* 9:553-563 (1993)) and, apoptosis is not observed in those animal models in which viral replication  
35 does not result in AIDS. *Id.* Further data indicates that uninfected but primed or activated T lymphocytes from HIV-infected individuals undergo apoptosis after

encountering the TNF-family ligand FasL. Using monocytic cell lines that result in death following HIV infection, it has been demonstrated that infection of U937 cells with HIV results in the *de novo* expression of FasL and that FasL mediates HIV-induced apoptosis (A.D. Badley *et al.*, *J. Virol.* 70:199-206 (1996)). Further, the  
5 TNF-family ligand was detectable in uninfected macrophages and its expression was upregulated following HIV infection resulting in selective killing of uninfected CD4 T-lymphocytes. *Id.* Thus, by the invention, a method for treating HIV<sup>+</sup> individuals is provided which involves administering an antagonist of the present invention to reduce selective killing of CD4 T-lymphocytes. Modes of administration and dosages are  
10 discussed in detail below.

In rejection of an allograft, the immune system of the recipient animal has not previously been primed to respond because the immune system for the most part is only primed by environmental antigens. Tissues from other members of the same species have not been presented in the same way that, for example, viruses and bacteria have  
15 been presented. In the case of allograft rejection, immunosuppressive regimens are designed to prevent the immune system from reaching the effector stage. However, the immune profile of xenograft rejection may resemble disease recurrence more than allograft rejection. In the case of disease recurrence, the immune system has already been activated, as evidenced by destruction of the native islet cells. Therefore, in  
20 disease recurrence, the immune system is already at the effector stage. Antagonists of the present invention are able to suppress the immune response to both allografts and xenografts because lymphocytes activated and differentiated into effector cells will express the TR10 polypeptide, and thereby are susceptible to compounds which enhance apoptosis. Thus, the present invention further provides a method for creating  
25 immune privileged tissues.

TR10 antagonists of the invention can further be used in the treatment of inflammatory diseases, such as inflammatory bowel disease, rheumatoid arthritis, osteoarthritis, psoriasis, and septicemia.

In addition, due to lymphoblast expression of TR10, soluble TR10 agonist or  
30 antagonist mABs may be used to treat this form of cancer.

### ***Modes of Administration***

The agonist or antagonists described herein can be administered *in vitro*, *ex vivo*, or *in vivo* to cells which express the receptor of the present invention. By  
35 administration of an "effective amount" of an agonist or antagonist is intended an amount of the compound that is sufficient to enhance or inhibit a cellular response to a

TNF-family ligand and include polypeptides. In particular, by administration of an "effective amount" of an agonist or antagonists is intended an amount effective to enhance or inhibit TR10 mediated apoptosis. Of course, where it is desired for apoptosis to be enhanced, an agonist according to the present invention can be co-administered with a TNF-family ligand. One of ordinary skill will appreciate that effective amounts of an agonist or antagonist can be determined empirically and may be employed in pure form or in pharmaceutically acceptable salt, ester or prodrug form. The agonist or antagonist may be administered in compositions in combination with one or more pharmaceutically acceptable excipients.

It will be understood that, when administered to a human patient, the total daily usage of the compounds and compositions of the present invention will be decided by the attending physician within the scope of sound medical judgment. The specific therapeutically effective dose level for any particular patient will depend upon factors well known in the medical arts.

As a general proposition, the total pharmaceutically effective amount of TR10 polypeptide administered parenterally per dose will be in the range of about 1 g/kg/day to 10 mg/kg/day of patient body weight, although, as noted above, this will be subject to therapeutic discretion. More preferably, this dose is at least 0.01 mg/kg/day, and most preferably for humans between about 0.01 and 1 mg/kg/day for the hormone. If given continuously, the TR10 polypeptide is typically administered at a dose rate of about 1 g/kg/hour to about 50 g/kg/hour, either by 1-4 injections per day or by continuous subcutaneous infusions, for example, using a mini-pump. An intravenous bag solution may also be employed.

Dosaging may also be arranged in a patient specific manner to provide a predetermined concentration of an agonist or antagonist in the blood, as determined by the RIA technique. Thus patient dosaging may be adjusted to achieve regular on-going trough blood levels, as measured by RIA, on the order of from 50 to 1000 ng/ml, preferably 150 to 500 ng/ml.

Pharmaceutical compositions containing the TR10 polypeptide of the invention may be administered orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, drops or transdermal patch), buccally, or as an oral or nasal spray. By "pharmaceutically acceptable carrier" is meant a non-toxic solid, semisolid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type. The term "parenteral" as used herein refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion.

Pharmaceutical compositions of the present invention for parenteral injection can comprise pharmaceutically acceptable sterile aqueous or nonaqueous solutions, dispersions, suspensions or emulsions as well as sterile powders for reconstitution into sterile injectable solutions or dispersions just prior to use.

5 In addition to soluble TR10 polypeptides, TR10 polypeptides containing the transmembrane region can also be used when appropriately solubilized by including detergents, such as CHAPS or NP-40, with buffer.

### *Chromosome assays*

10 The nucleic acid molecules of the present invention are also valuable for chromosome identification. The sequence is specifically targeted to and can hybridize with a particular location on an individual human chromosome. The mapping of DNAs to chromosomes according to the present invention is an important first step in correlating those sequences with genes associated with disease.

15 In certain preferred embodiments in this regard, the cDNA herein disclosed is used to clone genomic DNA of a TR10 receptor gene. This can be accomplished using a variety of well known techniques and libraries, which generally are available commercially. The genomic DNA is then used for *in situ* chromosome mapping using well known techniques for this purpose.

20 In addition, in some cases, sequences can be mapped to chromosomes by preparing PCR primers (preferably 15-25 bp) from the cDNA. Computer analysis of the 3' untranslated region of the gene is used to rapidly select primers that do not span more than one exon in the genomic DNA, thus complicating the amplification process. These primers are then used for PCR screening of somatic cell hybrids containing  
25 individual human chromosomes.

Fluorescence *in situ* hybridization ("FISH") of a cDNA clone to a metaphase chromosomal spread can be used to provide a precise chromosomal location in one step. This technique can be used with cDNA as short as 50 or 60 bp. For a review of this technique, see Verma *et al.*, *Human Chromosomes: a Manual of Basic Techniques*,  
30 Pergamon Press, New York (1988).

Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. Such data are found, for example, in V. McKusick, *Mendelian Inheritance in Man*, available on line through Johns Hopkins University, Welch Medical Library. The  
35 relationship between genes and diseases that have been mapped to the same

chromosomal region are then identified through linkage analysis (coinheritance of physically adjacent genes).

Next, it is necessary to determine the differences in the cDNA or genomic sequence between affected and unaffected individuals. If a mutation is observed in some or all of the affected individuals but not in any normal individuals, then the mutation is likely to be the causative agent of the disease.

Having generally described the invention, the same will be more readily understood by reference to the following examples, which are provided by way of illustration and are not intended as limiting.

### **Example 1**

#### ***Expression and Purification of the TR10 Receptor in E. coli***

The bacterial expression vector pQE60 is used for bacterial expression in this example. (QIAGEN, Inc., 9259 Eton Avenue, Chatsworth, CA, 91311). pQE60 encodes ampicillin antibiotic resistance ("Amp<sup>r</sup>") and contains a bacterial origin of replication ("ori"), an IPTG inducible promoter, a ribosome binding site ("RBS"), six codons encoding histidine residues that allow affinity purification using nickel-nitrilotri-acetic acid ("Ni-NTA") affinity resin sold by QIAGEN, Inc., *supra*, and suitable single restriction enzyme cleavage sites. These elements are arranged such that a DNA fragment encoding a polypeptide may be inserted in such a way as to produce that polypeptide with the six His residues (i.e., a "6 X His tag") covalently linked to the carboxyl terminus of that polypeptide. However, in this example, the polypeptide coding sequence is inserted such that translation of the six His codons is prevented and, therefore, the polypeptide is produced with no 6 X His tag.

The DNA sequence encoding the desired portion of the TR10 protein lacking the hydrophobic leader sequence is amplified from the deposited cDNA clone using PCR oligonucleotide primers which anneal to the amino terminal sequences of the desired portion of the TR10 protein and to sequences in the deposited construct 3' to the cDNA coding sequence. Additional nucleotides containing restriction sites to facilitate cloning in the pQE60 vector are added to the 5' and 3' sequences, respectively.

For cloning the mature protein, the 5' primer has the sequence:  
5'-CGCCCATGGCCACCATCCCCCGGCAG-3' (SEQ ID NO: 10) containing the underlined NcoI restriction site followed by nucleotides complementary to the amino terminal coding sequence of the mature TR10 sequence in Figure 1. One of ordinary skill in the art would appreciate, of course, that the point in the protein coding sequence where the 5' primer begins may be varied to amplify a desired portion of the complete protein shorter or longer than the mature form.



The 3' primer has the sequence:

5'-CGCAAGCTTTTAGTAGTGATAGGGAGAGGC-3' (SEQ ID NO:11) containing the underlined HindIII site followed by nucleotides complementary to the 3' end of the non-coding sequence in the TR10 DNA sequence in Figure 1.

5       The amplified TR10 DNA fragments and the vector pQE60 are digested with Nco I and HindIII and the digested DNAs then ligated together. Insertion of the TR10 protein DNA into the restricted pQE60 vector places the TR10 protein coding region (including its associated stop codon) downstream from the IPTG-inducible promoter and in-frame with an initiating AUG. The associated stop  
10       codon prevents translation of the six histidine codons downstream of the insertion point.

      The ligation mixture is transformed into competent *E. coli* cells using standard procedures. Such procedures are described in Sambrook *et al.*, *Molecular Cloning: a Laboratory Manual*, 2nd Ed.; Cold Spring Harbor Laboratory Press, Cold Spring  
15       Harbor, N.Y. (1989). *E. coli* strain M15/rep4, containing multiple copies of the plasmid pREP4, which expresses lac repressor and confers kanamycin resistance ("Kan"), is used in carrying out the illustrative example described herein. This strain, which is only one of many that are suitable for expressing TR10 protein, is available commercially from Qiagen, Inc., *supra*.

20       Transformants are identified by their ability to grow on LB plates in the presence of ampicillin and kanamycin. Plasmid DNA is isolated from resistant colonies and the identity of the cloned DNA confirmed by restriction analysis, PCR, and DNA sequencing.

      Clones containing the desired constructs are grown overnight ("O/N") in liquid  
25       culture in LB media supplemented with both ampicillin (100 g/ml) and kanamycin (25 g/ml). The O/N culture is used to inoculate a large culture, at a dilution of approximately 1:100 to 1:250. The cells are grown to an optical density at 600nm ("OD600") of between 0.4 and 0.6. Isopropyl-B-D-thiogalactopyranoside ("IPTG") is then added to a final concentration of 1 mM to induce transcription from the *lac*  
30       repressor sensitive promoter, by inactivating the *lacI* repressor. Cells subsequently are incubated further for 3 to 4 hours. Cells then are harvested by centrifugation.

      The cells are then stirred for 3-4 hours at 4°C in 6M guanidine-HCl, pH8. The cell debris is removed by centrifugation, and the supernatant containing the TR10 is loaded onto a nickel-nitrilo-tri-acetic acid ("NiNTA") affinity resin column (available  
35       from QIAGEN, Inc., *supra*). Proteins with a 6 x His tag bind to the NI-NTA resin with high affinity and can be purified in a simple one-step procedure (for details see:

The QIAexpressionist, 1995, QIAGEN, Inc., *supra*). Briefly the supernatant is loaded onto the column in 6 M guanidine-HCl, pH8, the column is first washed with 10 volumes of 6 M guanidine-HCl, pH8, then washed with 10 volumes of 6 M guanidine-HCl pH6, and finally the TR10 is eluted with 6 M guanidine-HCl, pH5.

5 The purified protein is then renatured by dialyzing it against phosphatebuffered saline (PBS) or 50 mM Na-acetate, pH 6 buffer plus 200 mM NaCl. Alternatively, the protein can be successfully refolded while immobilized on the Ni-NTA column. The recommended conditions are as follows: renature using a linear 6M-1M urea gradient in 500 mM NaCl, 20% glycerol, 20 mM Tris/HCl pH7.4, containing protease inhibitors.  
10 The renaturation should be performed over a period of 1.5 hours or more. After renaturation the proteins can be eluted by the addition of 250 mM imidazole. Imidazole is removed by a final dialyzing step against PBS or 50 mM sodium acetate pH6 buffer plus 200 mM NaCl. The purified protein is stored at 4°C or frozen at -80°C.

15

### **Example 2**

#### ***Cloning and Expression of TR10 in a Baculovirus Expression System***

In this illustrative example, the plasmid shuttle vector pA2 is used to insert the cloned DNA encoding the complete protein, including its naturally associated secretory  
20 signal (leader) sequence, into a baculovirus to express the mature TR10 protein, using standard methods as described in Summers *et al.*, *A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures*, Texas Agricultural Experimental Station Bulletin No. 1555 (1987). This expression vector contains the strong polyhedrin promoter of the *Autographa californica* nuclear polyhedrosis virus  
25 (AcMNPV) followed by convenient restriction sites such as BamHI and Asp718. The polyadenylation site of the simian virus 40 ("SV40") is used for efficient polyadenylation. For easy selection of recombinant virus, the plasmid contains the beta-galactosidase gene from *E. coli* under control of a weak *Drosophila* promoter in the same orientation, followed by the polyadenylation signal of the polyhedrin gene. The  
30 inserted genes are flanked on both sides by viral sequences for cell-mediated homologous recombination with wild-type viral DNA to generate viable virus that express the cloned polynucleotide.

Many other baculovirus vectors could be used in place of the vector above, such as pAc373, pVL941 and pAcIM1, as one skilled in the art would readily appreciate, as  
35 long as the construct provides appropriately located signals for transcription, translation, secretion and the like, including a signal peptide and an in-frame AUG as

required. Such vectors are described, for instance, in Luckow *et al.*, *Virology* 170:31-39 (1989).

The cDNA sequence encoding the mature TR10 receptor protein in the deposited clone, lacking the AUG initiation codon and the naturally associated leader sequence shown in Figure 1 (SEQ ID NO:2), is amplified using PCR oligonucleotide primers corresponding to the 5' and 3' sequences of the gene.

The 5' primer has the sequence 5' CGCGGATCC GCCATCATGGGACTTTGGGGACAA 3' (SEQ ID NO:12) containing the underlined BamHI restriction enzyme site, an efficient signal for initiation of translation in eukaryotic cells, as described by M. Kozak, *J. Mol. Biol.* 196:947-950 (1987), followed by bases of the sequence of the mature TR10 protein shown in Figure 1, beginning with the indicated N-terminus of the mature protein.

The 3' primer for TR10 has the sequence 5' CGCGGTACCTTAGTAGTGATAGGGAGAGGC 3' (SEQ ID NO:13) containing the underlined Asp718 restriction site followed by nucleotides complementary to the 3' noncoding sequence in Figure 1.

The amplified fragment is isolated from a 1% agarose gel using a commercially available kit ("GeneClean," BIO 101 Inc., La Jolla, Ca.) The fragment then is digested with BamHI and Asp718 and again is purified on a 1% agarose gel. This fragment is designated "F1."

The plasmid is digested with the restriction enzyme Bam HI and optionally can be dephosphorylated using calf intestinal phosphatase, using routine procedures known in the art. The DNA is then isolated from a 1% agarose gel using a commercially available kit ("GeneClean" BIO 101 Inc., La Jolla, Ca.). The vector DNA is designated herein "V1."

Fragment F1 and the dephosphorylated plasmid V1 are ligated together with T4 DNA ligase. *E. coli* HB101 or other suitable *E. coli* hosts such as XL-1 Blue (Stratagene Cloning Systems, La Jolla, CA) cells are transformed with the ligation mixture and spread on culture plates. Bacteria are identified that contain the plasmid with the human TR10 gene using the PCR method, in which one of the primers that is used to amplify the gene and the second primer is from well within the vector so that only those bacterial colonies containing the TR10 gene fragment will show amplification of the DNA. The sequence of the cloned fragment is confirmed by DNA sequencing. This plasmid is designated herein pBacTR10.

Five  $\mu$ g of the plasmid pBacTR10 is co-transfected with 1.0  $\mu$ g of a commercially available linearized baculovirus DNA ("BaculoGold™ baculovirus

DNA", Pharmingen, San Diego, CA.), using the lipofectin method described by Felgner *et al.*, *Proc. Natl. Acad. Sci. USA* 84:7413-7417 (1987). 1  $\mu$ g of BaculoGold™ virus DNA and 5  $\mu$ g of the plasmid pBacTR10 are mixed in a sterile well of a microliter plate containing 50  $\mu$ l of serum free Grace's medium (Life Technologies, Inc., Rockville, MD). Afterwards, 10  $\mu$ l Lipofectin plus 90  $\mu$ l Grace's medium are added, mixed, and incubated for 15 minutes at room temperature. Then, the transfection mixture is added drop-wise to Sf9 insect cells (ATCC CRL 1711) seeded in a 35 mm tissue culture plate with 1 ml Grace's medium without serum. The plate is rocked back and forth to mix the newly added solution. The plate is then incubated for 5 hours at 27°C. After 5 hours, the transfection solution is removed from the plate and 1 ml of Grace's insect medium supplemented with 10% fetal calf serum is added. The plate is put back into an incubator and cultivation is continued at 27°C for four days.

After four days, the supernatant is collected and a plaque assay is performed, as described by Summers and Smith, cited above. An agarose gel with "Blue Gal" (Life Technologies, Inc., Rockville, MD) is used to allow easy identification and isolation of gal-expressing clones, which produce blue-stained plaques. (A detailed description of a "plaque assay" of this type can also be found in the user's guide for insect cell culture and baculovirology distributed by Life Technologies, Inc., Rockville, MD, pages 9-10). After appropriate incubation, blue stained plaques are picked with the tip of a micropipettor (e.g., Eppendorf). The agar containing the recombinant viruses is then resuspended in a microcentrifuge tube containing 200  $\mu$ l of Grace's medium and the suspension containing the recombinant baculovirus is used to infect Sf9 cells seeded in 35 mm dishes. Four days later the supernatants of these culture dishes are harvested and then they are stored at 4°C. The recombinant virus is called V-TR10.

To verify the expression of the TR10 gene, Sf9 cells are grown in Grace's medium supplemented with 10% heat inactivated FBS. The cells are infected with the recombinant baculovirus V-TR10 at a multiplicity of infection ("MOI") of about 2. Six hours later the medium is removed and is replaced with SF900 II medium minus methionine and cysteine (available from Life Technologies, Inc., Rockville, MD). If radiolabeled proteins are desired, 42 hours later, 5  $\mu$ Ci of  $^{35}$ S-methionine and 5  $\mu$ Ci  $^{35}$ S-cysteine (available from Amersham) are added. The cells are further incubated for 16 hours and then they are harvested by centrifugation. The proteins in the supernatant as well as the intracellular proteins are analyzed by SDS-PAGE followed by autoradiography (if radiolabeled). Microsequencing of the amino acid sequence of the amino terminus of purified protein may be used to determine the amino terminal

sequence of the mature protein and thus the cleavage point and length of the secretory signal peptide.

### **Example 3**

#### **5 Cloning and Expression of the TR10 Receptor in Mammalian Cells**

A typical mammalian expression vector contains the promoter element, which mediates the initiation of transcription of mRNA, the protein coding sequence, and signals required for the termination of transcription and polyadenylation of the transcript. Additional elements include enhancers, Kozak sequences and intervening sequences flanked by donor and acceptor sites for RNA splicing. Highly efficient transcription can be achieved with the early and late promoters from SV40, the long terminal repeats (LTRs) from Retroviruses, e.g. RSV, HTLVI, HIVI and the early promoter of the cytomegalovirus (CMV). However, cellular signals can also be used (e.g., the human actin promoter). Suitable expression vectors for use in practicing the present invention include, for example, vectors such as pSVL and pMSG (Pharmacia, Uppsala, Sweden), pRSVcat (ATCC 37152), pSV2dhfr (ATCC 37146) and pBC12MI (ATCC 67109). Mammalian host cells that could be used include, human Hela 293, H9 and Jurkat cells, mouse NIH3T3 and C127 cells, Cos 1, Cos 7 and CV1, quail QC1-3 cells, mouse L cells, and Chinese hamster ovary (CHO) cells.

Alternatively, the gene can be expressed in stable cell lines that contain the gene integrated into a chromosome. Co-transfection with a selectable marker such as dhfr, gpt, neomycin, or hygromycin allows the identification and isolation of the transfected cells.

The transfected gene can also be amplified to express large amounts of the encoded protein. The dihydrofolate reductase (DHFR) marker is useful to develop cell lines that carry several hundred or even several thousand copies of the gene of interest. Another useful selection marker is the enzyme glutamine synthase (GS) (Murphy *et al.*, *Biochem. J.* 227:277-279 (1991); Bebbington *et al.*, *Bio/Technology* 10:169-175 (1992)). Using these markers, the mammalian cells are grown in selective medium and the cells with the highest resistance selected. These cell lines contain the amplified gene(s) integrated into a chromosome. Chinese hamster ovary (CHO) cells are often used for the production of proteins.

The expression vectors pCl and pC4 contain the strong promoter (LTR) of the Rous Sarcoma Virus (Cullen *et al.*, *Molecular and Cellular Biology* 5:438-447 (March 1985)), plus a fragment of the CMV-enhancer (Boshart *et al.*, *Cell* 41:521-530 (1985)). Multiple cloning sites, e.g., with the restriction enzyme cleavage sites BamHI, XbaI

and Asp718, facilitate the cloning of the gene of interest. The vectors contain in addition the 3' intron, the polyadenylation and termination signal of the rat preproinsulin gene.

**Example 3A**

**5 Cloning and Expression of the Extracellular Soluble Domain of TR10 in COS cells**

The expression plasmid, pTR10-HA, is made by cloning a cDNA encoding TR10 into the expression vector pcDNA1/Amp or pcDNAIII (which can be obtained from Invitrogen, Inc.).

10 The expression vector pcDNA1/amp contains: (1) an *E. coli* origin of replication effective for propagation in *E. coli* and other prokaryotic cell; (2) an ampicillin resistance gene for selection of plasmid-containing prokaryotic cells; (3) an SV40 origin of replication for propagation in eukaryotic cells; (4) a CMV promoter, a polylinker, an SV40 intron, and a polyadenylation signal arranged so that a cDNA conveniently can be  
15 placed under expression control of the CMV promoter and operably linked to the SV40 intron and the polyadenylation signal by means of restriction sites in the polylinker.

A DNA fragment encoding the entire TR10 precursor and a HA tag fused in frame to its 3' end is cloned into the polylinker region of the vector so that recombinant protein expression is directed by the CMV promoter. The HA tag corresponds to an  
20 epitope derived from the influenza hemagglutinin protein described by Wilson *et al.*, *Cell* 37:767 (1984). The fusion of the HA tag to the target protein allows easy detection of the recombinant protein with an antibody that recognizes the HA epitope.

The plasmid construction strategy is as follows:

The TR10 cDNA of the deposited clone is amplified using primers that contain  
25 convenient restriction sites, much as described above regarding the construction of expression vectors for expression of TR10 in *E. coli*.

To facilitate detection, purification and characterization of the expressed TR10, one of the primers contains a hemagglutinin tag ("HA tag") as described above.

Suitable primers for TR10 include the following, which are used in this  
30 example:

The 5' primer, 5' CGCGGATCCGCCATCATGGGACTTTGGGGACAA 3' (SEQ ID NO:12) contains the underlined BamHI site, an ATG start codon and 5 codons thereafter. The 3' primer for TR10, which contains the underlined XbaI site, stop codon, hemagglutinin tag, and the last 19 nucleotides of the 3' coding sequence (at the  
35 3' end), has the following sequence:

5'CGCTCTAGATCAAGCGTAGTCTGGGACGTCGTATGGGTAGTAAGT  
GATAGGGAGAGGC 3' (SEQ ID NO:14).



The PCR amplified DNA fragment and the vector, pcDNAI/Amp, are digested with BamHI and XbaI and then ligated. The ligation mixture is transformed into *E. coli* strain SURE (available from Stratagene Cloning Systems, 11099 North Torrey Pines Road, La Jolla, CA 92037) the transformed culture is plated on ampicillin media plates which then are incubated to allow growth of ampicillin resistant colonies. Plasmid DNA is isolated from resistant colonies and examined by restriction analysis and gel sizing for the presence of the TR10-encoding fragment.

For expression of recombinant TR10, COS cells are transfected with an expression vector, as described above, using DEAE-DEXTRAN, as described, for instance, in Sambrook *et al.*, *Molecular Cloning: a Laboratory Manual*, Cold Spring Laboratory Press, Cold Spring Harbor, NY (1989). Cells are incubated under conditions for expression of TR10 by the vector.

Expression of the TR10-HA fusion protein is detected by radiolabelling and immunoprecipitation, using methods described in, for example Harlow *et al.*, *Antibodies: a Laboratory Manual*, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY (1988). To this end, two days after transfection, the cells are labeled by incubation in media containing <sup>35</sup>S-cysteine for 8 hours. The cells and the media are collected, and the cells are washed and then lysed with detergent-containing RIPA buffer: 150 mM NaCl, 1% NP-40, 0.1% SDS, 1% NP-40, 0.5% DOC, 50 mM TRIS, pH 7.5, as described by Wilson *et al.* cited above. Proteins are precipitated from the cell lysate and from the culture media using an HA-specific monoclonal antibody. The precipitated proteins then are analyzed by SDS-PAGE gels and autoradiography. An expression product of the expected size is seen in the cell lysate, which is not seen in negative controls.

### **Example 3B**

#### ***Cloning and Expression of TR10 using the CHO Expression System***

The vector pC4 is used for the expression of the TR10 polypeptide. Plasmid pC4 is a derivative of the plasmid pSV2-dhfr (ATCC Accession No. 37146). The plasmid contains the mouse DHFR gene under control of the SV40 early promoter. Chinese hamster ovary- or other cells lacking dihydrofolate activity that are transfected with these plasmids can be selected by growing the cells in a selective medium (alpha minus MEM, Life Technologies, Rockville, MD) supplemented with the chemotherapeutic agent methotrexate (MTX). The amplification of the DHFR genes in cells resistant to MTX has been well documented (see, e.g., F.W. Alt *et al.*, *J. Biol. Chem.* 253:1357-1370 (1978); J.L. Hamlin and C. Ma, *Biochem. et Biophys. Acta* 1097:107-143 (1990); M.J. Page M.A. Sydenham, *Biotechnology* 9:64-68(1991)).

Cells grown in increasing concentrations of MTX develop resistance to the drug by overproducing the target enzyme, DHFR, as a result of amplification of the DHFR gene. If a second gene is linked to the DHFR gene, it is usually co-amplified and over-expressed. It is known in the art that this approach may be used to develop cell lines  
5 carrying more than 1,000 copies of the amplified gene(s). Subsequently, when the methotrexate is withdrawn, cell lines are obtained that contain the amplified gene integrated into one or more chromosome(s) of the host cell.

Plasmid pC4 contains, for expressing the gene of interest, the strong promoter of the long terminal repeat (LTR) of the Rous Sarcoma Virus (Cullen *et al.*, *Molecular and Cellular Biology* 5:438-447 (March 1985)), plus a fragment isolated from the  
10 enhancer of the immediate early gene of human cytomegalovirus (CMV) (Boshart *et al.*, *Cell* 41:521-530 (1985)). Downstream of the promoter are the following single restriction enzyme cleavage sites that allow the integration of the genes: BamHI, XbaI, and Asp718. Behind these cloning sites, the plasmid contains the 3' intron and the  
15 polyadenylation site of the rat preproinsulin gene. Other high efficiency promoters can also be used for the expression, e.g., the human B-actin promoter, the SV40 early or late promoters or the long terminal repeats from other retroviruses, e.g., HIV and HTLVI. Clontech's Tet-Off and Tet-On gene expression systems and similar systems can be used to express the TR10 polypeptide in a regulated way in mammalian cells.  
20 For the polyadenylation of the mRNA, other signals, e.g., from the human growth hormone or globin genes, can be used as well.

Stable cell lines carrying a gene of interest integrated into the chromosomes can also be selected upon co-transfection with a selectable marker such as gpt, G418, or hygromycin. It is advantageous to use more than one selectable marker in the  
25 beginning, e.g., G418 plus methotrexate.

The plasmid pC4 is digested with the restriction enzyme BamHI and then dephosphorylated using calf intestinal phosphates, by procedures known in the art. The vector is then isolated from a 1% agarose gel.

The DNA sequence encoding the complete TR10 polypeptide is amplified using  
30 PCR oligonucleotide primers corresponding to the 5' and 3' sequences of the desired portion of the gene.

The 5' oligonucleotide primer for TR10, containing the underlined BamHI restriction site, a Kozak sequence, and an AUG start codon, has the sequence:  
5' CGCGGATCCGCCATCATGGGACTTTGGGGACAA 3' (SEQ ID NO:12). The  
35 3' primer for TR10, containing the underlined Asp718 restriction site, has the sequence:  
5' CGCGGTACCTTAGTAGTGATAGGGAGAGGC 3' (SEQ ID NO:13).

The amplified fragment is digested with BamHI and then purified again on a 1% agarose gel. The isolated fragment and the dephosphorylated vector are then ligated with T4 DNA ligase. E. coli HB 101 or XL-1 Blue cells are then transformed and bacteria are identified that contain the fragment inserted into plasmid pC4 using, for instance, restriction enzyme analysis.

Chinese hamster ovary cells lacking an active DHFR enzyme are used for transfection. Five  $\mu$ g of the expression plasmid pC4 are cotransfected with 0.5  $\mu$ g of the plasmid pSVneo using the lipofectin method (Felgner *et al.*, *supra*). The plasmid pSV2-neo contains a dominant selectable marker, the *neo* gene from Tn5 encoding an enzyme that confers resistance to a group of antibiotics including G418. The cells are seeded in alpha minus MEM supplemented with 1 mg/ml G418. After 2 days, the cells are trypsinized and seeded in hybridoma cloning plates (Greiner, Germany) in alpha minus MEM supplemented with 10, 25, or 50 ng/ml of MTX plus 1 mg/ml G418. After about 10-14 days, single clones are trypsinized and then seeded in 6-well petri dishes or 10 ml flasks using different concentrations of methotrexate (50 nM, 100 nM, 200 nM, 400 nM, 800nM). Clones growing at the highest concentrations of methotrexate are then transferred to new 6-well plates containing even higher concentrations of methotrexate (1  $\mu$ M, 2  $\mu$ M, 5  $\mu$ M, 10  $\mu$ M, 20  $\mu$ M). The same procedure is repeated until clones are obtained which grow at a concentration of 100-200  $\mu$ M. Expression of the desired gene product is analyzed, for instance, by Western blot analysis and SDS-PAGE, or by reversed phase HPLC analysis.

#### **Example 4**

##### ***Tissue distribution of TR10 mRNA expression***

Northern blot analysis was carried out to examine TR10 gene expression in human tissues, using methods described by, among others, Sambrook *et al.*, cited above. A cDNA probe containing the entire nucleotide sequence of the TR10 protein (SEQ ID NO:1) was labeled with  $^{32}$ P using the *rediprime*<sup>TM</sup> DNA labeling system (Amersham Life Science), according to manufacturer's instructions. After labeling, the probe was purified using a CHROMA SPIN-100 column (Clontech Laboratories, Inc.), according to manufacturer's protocol number PT1200-1. The purified labeled probe was then used to examine various human tissues for TR10 mRNA.

Multiple Tissue Northern (MTN) blots containing various human tissues (H) or human immune system tissues (IM) were obtained from Clontech and were examined with labeled probe using ExpressHyb<sup>TM</sup> hybridization solution (Clontech) according to manufacturer's protocol number PT1190-1. Following hybridization and washing, the blots were mounted and exposed to film at -70°C overnight, and films developed

according to standard procedures. Expression of TR10 was detected in tissues enriched in lymphocytes including peripheral blood leukocytes (PBLs), fetal liver, lung, kidney, small intestine, colon, keratinocytes, endothelial cells, and monocyte activated tissue. It can be envisaged that TR10 plays a role in lymphocyte homeostasis.

5

### *Northern Blot analysis of TR10 in various cell lines*

#### *Methods*

##### *Cells*

Unless stated otherwise, cell lines were obtained from the American Type  
10 Culture Collection (Rockville, MD). The myeloid (Koeffler *et al.* (1980); Koeffler  
(1983); Harris and Ralph (1985); and Tucker *et al.* (1987) and B-cell lines (Jonak *et al.*  
(1922)) studied represent cell types at different stages of the differentiation pathway.  
KG1a and PLB 985 cells (Tucker *et al.* (1987)) were obtained from H.P. Koeffler  
(UCLA School of Medicine). BJA-B was from Z. Jonak (SmithKline Beecham).  
15 TF274, a stromal cell line exhibiting osteoblastic features, was generated from the bone  
marrow of a healthy male donor (Z. Jonak and K.B. Tan, unpublished). Primary  
carotid artery endothelial cells were purchased from Clonetics Corp. (San Diego, CA)  
and monocytes were prepared by differential centrifugation of peripheral blood  
mononuclear cells and adhesion to tissue culture dish. CD19+, CD4+ and CD8+ cells  
20 (>90% pure) were isolated with cell type specific immunomagnetic beads (Dynal, Lake  
Success, NY).

##### *RNA Analysis*

Total RNA of adult tissues were purchased from Clontech (Palo Alto, CA).  
25 Total RNA was extracted from cell lines (in exponential growth phase) and primary  
cells with TriReagent (Molecular Research Center, Inc., Cincinnati, OH). 5 to 7.5 g  
of total RNA was fractionated in a 1% agarose gel containing formaldehyde cast in a  
Wide Mini-Sub Cell gel tray (Bio-Rad, Hercules, CA) as described (Sambrook, et al.)  
with slight modifications. The formaldehyde concentration was reduced to 0.5M and  
30 the RNA was stained prior to electrophoresis with 100 g/ml of etidium bromide that  
was added to the loading buffer. After electrophoresis with continuous buffer  
recirculation (60 volts/90 min), the gel was photographed and the RNA was transferred  
quantitatively to Zeta-probe nylon membrane (Biorad, Hercules, CA) by vacuum-  
blotting with 25 mM NaOH for 90 min. After neutralization for 5-10 min, with 1M  
35 Tris-HCl, pH 7.5 containing 3M NaCl, the blots were prehybridized with 50%  
formamide, 8% dextran sulfate, 6xSSPE, 0.1% SDS and 100 g/ml of sheared and

denatured salmon sperm DNA for at least 30 min at 42°C. cDNA inserts labeled with <sup>32</sup>P-dCTP by random priming (Stratagene, La Jolla, CA), were denatured with 0.25M NaOH (10 min at 37°C) and added to the prehybridization solution. After 24-65 hr at 42°C, the blots were washed under high stringency conditions (Sambrook, *et al.*) and  
5 exposed to X-ray films.

### **Results**

Expression of TR10 was assessed by Northern blot in the following cell lines: HL60 (promyelocytic leukemia), Hela cell S3, K562 (chronic myelogenous leukemia),  
10 MOLT4 (lymphoblast leukemia), Raji (Burkitt's lymphoma), SW480 (colorectal adenocarcinoma), A549 (lung carcinoma), and G361 (melanoma), and could only be detected in Hela cell S3, SW480 (colorectal adenocarcinoma), and the A549 (lung carcinoma) cell lines.

### **Example 5**

#### **TR10 Induced Apoptosis**

Since the entire sequence of TR10, especially its extracellular cysteine-rich domains, is highly homologous to that of other TRAIL receptors, the ability to TRAIL to bind TR10 and induce apoptosis was assessed.  
20

#### **Experimental Design**

To facilitate detection, TR10 (amino acids 56-386) was cloned into pCMV2FLAG (IBI Kodak) as an in-frame fusion to the signal sequence and FLAG-epitope tag encoded by the vector. The cDNA encoding the extracellular domain of  
25 TR10 (amino acids 56-210) was obtained by PCR, similar to the methods described above, and subcloned into a modified pCMV1FLAG vector that allowed for in-frame fusion with the Fc portion of human IgG. DR4-Fc, TNRF1-Fc, Fc and soluble TRAIL and TNF alpha expression constructs have been described previously, Pan, G. et al., Science 276:111-113 (1997), which is incorporated herein by reference in its entirety.

30 The receptor-Fc fusions and soluble ligands were prepared and in vivo binding was performed as previously described, Pan G. et al., Science 276:111-113 (1997), and Pan G. et al., Science 277:815-818 (1997), both of which are incorporated herein by reference in their entirety.

Cell death blocking assays using receptor-Fc fusions were carried out as  
35 described previously by Pan G. et al., Science 276:111-113 (1997), and Pan G. et al.,

Science 277:815-818 (1997), both of which are incorporated herein by reference in their entirety.

### **Results**

5       The extracellular domain of TR10 was expressed as a secreted chimera fused to the Fc portion of human IgG in 293 cells. Conditioned medium from transfected cells was mixed with bacterially expressed soluble His-FLAG-tagged TRAIL. The resulting complex was precipitated with protein G-Sepharose and bound TRAIL detected by Western blotting with anti-FLAG antibody. Like DR4, DR5, and TR5 (TRID), TR10  
10 bound TRAIL. Corroborating this ability to bind TRAIL was the finding that TR10-Fc, like DR4-Fc, could efficiently block TRAIL-induced apoptosis.

      In keeping with TR10 possessing a truncated non-functional death domain was the observation that TR10 overexpression did not cause cell death in Hela cells, and as might be expected, could act as a dominant negative receptor antagonizing TRAIL-  
15 induced apoptosis. Therefore, ectopic expression of TR10, like that of the decoy receptor TR5, is capable of substantially attenuating TRAIL-induced cell death, suggesting that TR10 antagonizes TRAIL signaling.

      It will be clear that the invention may be practiced otherwise than as particularly described in the foregoing description and examples.

20       Numerous modifications and variations of the present invention are possible in light of the above teachings and, therefore, are within the scope of the appended claims.

      The entire disclosure of all publications (including patents, patent applications, journal articles, laboratory manuals, books, or other documents) cited herein are hereby incorporated by reference.



## Sequence Listing

### (1) GENERAL INFORMATION:

- (i) APPLICANTS: NI, et al.
- (ii) TITLE OF INVENTION: HUMAN TUMOR NECROSIS FACTOR RECEPTOR TR10
- (iii) NUMBER OF SEQUENCES: 15
- (iv) CORRESPONDENCE ADDRESS:
  - (A) ADDRESSEE: HUMAN GENOME SCIENCES, INC.
  - (B) STREET: 9410 KEY WEST AVENUE
  - (C) CITY: ROCKVILLE
  - (D) STATE: MD
  - (E) COUNTRY: US
  - (F) ZIP: 20850
- (v) COMPUTER READABLE FORM:
  - (A) MEDIUM TYPE: Floppy disk
  - (B) COMPUTER: IBM PC compatible
  - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
  - (D) SOFTWARE: PatentIn Release #1.0, Version #1.30
- (vi) CURRENT APPLICATION DATA:
  - (A) APPLICATION NUMBER:
  - (B) FILING DATE:
  - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
  - (A) NAME: BROOKES, ANDERS A.
  - (B) REGISTRATION NUMBER: 36,373
  - (C) REFERENCE/DOCKET NUMBER: PF379
- (ix) TELECOMMUNICATION INFORMATION:
  - (A) TELEPHONE: (301) 309-8504
  - (B) TELEFAX: (301) 309-8512

### (2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 3566 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- (ix) FEATURE:
  - (A) NAME/KEY: CDS
  - (B) LOCATION: 109..1266
- (ix) FEATURE:
  - (A) NAME/KEY: sig\_peptide
  - (B) LOCATION: 109..271

## (ix) FEATURE:

(A) NAME/KEY: mat\_peptide

(B) LOCATION: 274..1266

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

CGACCCACGC GTCCGCCAC GCGTCCGGAG AACCTTTGCA CGCGCACAAA CTACGGGGAC	60
GATTTCTGAT TGATTTTGG CGCTTTCGAT CCACCTCCT CCCTTCTC ATG GGA CTT	117
Met Gly Leu	
-55	
TGG GGA CAA AGC GTC CCG ACC GCC TCG AGC GCT CGA GCA GGG CGC TAT	165
Trp Gly Gln Ser Val Pro Thr Ala Ser Ser Ala Arg Ala Gly Arg Tyr	
-50 -45 -40	
CCA GGA GCC AGG ACA GCG TCG GGA ACC AGA CCA TGG CTC CTG GAC CCC	213
Pro Gly Ala Arg Thr Ala Ser Gly Thr Arg Pro Trp Leu Leu Asp Pro	
-35 -30 -25	
AAG ATC CTT AAG TTC GTC GTC TTC ATC GTC GCG GTT CTG CTG CCG GTC	261
Lys Ile Leu Lys Phe Val Val Phe Ile Val Ala Val Leu Leu Pro Val	
-20 -15 -10 -5	
CGG GTT GAC TCT GCC ACC ATC CCC CGG CAG GAC GAA GTT CCC CAG CAG	309
Arg Val Asp Ser Ala Thr Ile Pro Arg Gln Asp Glu Val Pro Gln Gln	
1 5 10	
ACA GTG GCC CCA CAG CAA CAG AGG CGC AGC CTC AAG GAG GAG GAG TGT	357
Thr Val Ala Pro Gln Gln Gln Arg Arg Ser Leu Lys Glu Glu Glu Cys	
15 20 25	
CCA GCA GGA TCT CAT AGA TCA GAA TAT ACT GGA GCC TGT AAC CCG TGC	405
Pro Ala Gly Ser His Arg Ser Glu Tyr Thr Gly Ala Cys Asn Pro Cys	
30 35 40	
ACA GAG GGT GTG GAT TAC ACC ATT GCT TCC AAC AAT TTG CCT TCT TGC	453
Thr Glu Gly Val Asp Tyr Thr Ile Ala Ser Asn Asn Leu Pro Ser Cys	
45 50 55 60	
CTG CTA TGT ACA GTT TGT AAA TCA GGT CAA ACA AAT AAA AGT TCC TGT	501
Leu Leu Cys Thr Val Cys Lys Ser Gly Gln Thr Asn Lys Ser Ser Cys	
65 70 75	
ACC ACG ACC AGA GAC ACC GTG TGT CAG TGT GAA AAA GGA AGC TTC CAG	549
Thr Thr Thr Arg Asp Thr Val Cys Gln Cys Glu Lys Gly Ser Phe Gln	
80 85 90	
GAT AAA AAC TCC CCT GAG ATG TGC CGG ACG TGT AGA ACA GGG TGT CCC	597
Asp Lys Asn Ser Pro Glu Met Cys Arg Thr Cys Arg Thr Gly Cys Pro	
95 100 105	
AGA GGG ATG GTC AAG GTC AGT AAT TGT ACG CCC CGG AGT GAC ATC AAG	645
Arg Gly Met Val Lys Val Ser Asn Cys Thr Pro Arg Ser Asp Ile Lys	
110 115 120	
TGC AAA AAT GAA TCA GCT GCC AGT TCC ACT GGG AAA ACC CCA GCA GCG	693
Cys Lys Asn Glu Ser Ala Ala Ser Ser Thr Gly Lys Thr Pro Ala Ala	
125 130 135 140	

GAG GAG ACA GTG ACC ACC ATC CTG GGG ATG CTT GCC TCT CCC TAT CAC Glu Glu Thr Val Thr Thr Ile Leu Gly Met Leu Ala Ser Pro Tyr His 145 150 155	741
TAC CTT ATC ATC ATA GTG GTT TTA GTC ATC ATT TTA GCT GTG GTT GTG Tyr Leu Ile Ile Ile Val Val Leu Val Ile Ile Leu Ala Val Val Val 160 165 170	789
GTT GGC TTT TCA TGT CGG AAG AAA TTC ATT TCT TAC CTC AAA GGC ATC Val Gly Phe Ser Cys Arg Lys Lys Phe Ile Ser Tyr Leu Lys Gly Ile 175 180 185	837
TGC TCA GGT GGT GGA GGA GGT CCC GAA CGT GTG CAC AGA GTC CTT TTC Cys Ser Gly Gly Gly Gly Gly Pro Glu Arg Val His Arg Val Leu Phe 190 195 200	885
CGG CGG CGT TCA TGT CCT TCA CGA GGT CCT GGG GCG GAG GAC AAT GCC Arg Arg Arg Ser Cys Pro Ser Arg Val Pro Gly Ala Glu Asp Asn Ala 205 210 215 220	933
CGC AAC GAG ACC CTG AGT AAC AGA TAC TTG CAG CCC ACC CAG GTC TCT Arg Asn Glu Thr Leu Ser Asn Arg Tyr Leu Gln Pro Thr Gln Val Ser 225 230 235	981
GAG CAG GAA ATC CAA GGT CAG GAG CTG GCA GAG CTA ACA GGT GTG ACT Glu Gln Glu Ile Gln Gly Gln Glu Leu Ala Glu Leu Thr Gly Val Thr 240 245 250	1029
GTA GAG TCG CCA GAG GAG CCA CAG CGT CTG CTG GAA CAG GCA GAA GCT Val Glu Ser Pro Glu Glu Pro Gln Arg Leu Leu Glu Gln Ala Glu Ala 255 260 265	1077
GAA GGG TGT CAG AGG AGG AGG CTG CTG GTT CCA GTG AAT GAC GCT GAC Glu Gly Cys Gln Arg Arg Arg Leu Leu Val Pro Val Asn Asp Ala Asp 270 275 280	1125
TCC GCT GAC ATC AGC ACC TTG CTG GAT GCC TCG GCA ACA CTG GAA GAA Ser Ala Asp Ile Ser Thr Leu Leu Asp Ala Ser Ala Thr Leu Glu Glu 285 290 295 300	1173
GGA CAT GCA AAG GAA ACA ATT CAG GAC CAA CTG GTG GGC TCC GAA AAG Gly His Ala Lys Glu Thr Ile Gln Asp Gln Leu Val Gly Ser Glu Lys 305 310 315	1221
CTC TTT TAT GAA GAA GAT GAG GCA GGC TCT GCT ACG TCC TGC CTG Leu Phe Tyr Glu Glu Asp Glu Ala Gly Ser Ala Thr Ser Cys Leu 320 325 330	1266
TGAAAGAATC TCTTCAGGAA ACCAGAGCTT CCTTCATTTA CCTTTTCTCC TACAAAGGGA	1326
AGCAGCCTGG AAGAAACAGT CCAGTACTTG ACCCATGCCC CAACAAACTC TACTATCCAA	1386
TATGGGGCAG CTTACCAATG GTCCTAGAAC TTTGTTAACG CACTTGGAGT AATTTTATG	1446
AAATACTGCG TGTGATAAGC AAACGGGAGA AATTTATATC AGATTCTTGG CTGCATAGTT	1506
ATACGATTGT GTATTAAGGG TCGTTTTAGG CCACATGCGG TGGCTCATGC CTGTAATCCC	1566
AGCACTTTGA TAGGCTGAGG CAGGTGGATT GCTTTGAGCT CGGGAGTTTG AGACCAGCCT	1626

CATCAACACA GTGAAACTCC ATCTCAATTT AAAAAGAAAA AAAAGTGGTT TTAGGATGTC	1686
ATTCTTTGCA GTTCTTCATC ATGAGACAAG TCTTTTTTTC TGCTTCTTAT ATTGCAAGCT	1746
CCATCTCTAC TGGTGTGTGC ATTTAATGAC ATCTAACTAC AGATGCCGCA CAGCCACAAT	1806
GCTTTGCCTT ATAGTTTTTT AACTTTAGAA CGGGATTATC TTGTTATTAC CTGTATTTTC	1866
AGTTTCGGAT ATTTTGA CT TAATGATGAG ATTATCAAGA CGTAGCCCTA TGCTAAGTCA	1926
TGAGCATATG GACTTACGAG GGTTCGACTT AGAGTTTTGA GCTTTAAGAT AGGATTATTG	1986
GGGCTTACCC CCACCTTAAT TAGAGAAACA TTTATATTGC TTA CTACTGT AGGCTGTACA	2046
TCTCTTTTCC GATTTTTGTA TAATGATGTA AACATGGAAA AACTTTAGGA AATGCACTTA	2106
TTAGGCTGTT TACATGGGTT GCCTGGATAC AAATCAGCAG TCAAAAATGA CTAAAAATAT	2166
AACTAGTGAC GGAGGGAGAA ATCCTCCCTC TGTGGGAGGC ACTTACTGCA TTCCAGTTCT	2226
CCCTCCTGCG CCCTGAGACT GGACCAGGGT TTGATGGCTG GCAGCTTCTC AAGGGGCAGC	2286
TTGCTTACT TGTAAATTTT AGAGGTATAT AGCCATATTT ATTTATAAAT AAATATTTAT	2346
TTATTTATTT ATAAGTAGAT GTTTACATAT GCCCAGGATT TTGAAGAGCC TGGTATCTTT	2406
GGGAAGCCAT GTGTCTGGTT TGTCGTGCTG GGACAGTCAT GGGACTGCAT CTTCCGACTT	2466
GTCCACAGCA GATGAGGACA GTGAGAATTA AGTTAGATCC GAGACTGCGA AGAGCTTCTC	2526
TTTCAAGCGC CATTACAGTT GAACGTTAGT GAATCTTGAG CCTCATTTGG GCTCAGGGCA	2586
GAGCAGGTGT TTATCTGCCC CGGCATCTGC CATGGCATCA AGAGGGAAGA GTGGACGGTG	2646
CTTGGGAATG GTGTGAAATG GTTGCCGACT CAGGCATGGA TGGGCCCCCTC TCGCTTCTGG	2706
TGGTCTGTGA ACTGAGTCCC TGGGATGCCT TTTAGGGCAG AGATTCCTGA GCTGCGTTTT	2766
AGGGTACAGA TTCCCTGTTT GAGGAGCTTG GCCCTCTGT AAGCATCTGA CTCATCTCAG	2826
AGATATCAAT TCTTAAACAC TGTGACAACG GGATCTAAAA TGGCTGACAC ATTTGTCCCTT	2886
GTGTCACGTT CCATTATTTT ATTTAAAAAC CTCAGTAATC GTTTTAGCTT CTTTCCAGCA	2946
AACTCTTCTC CACAGTAGCC CAGTCGTGGT AGGATAAATT ACGGATATAG TCATTCTAGG	3006
GGTTTCAGTC TTTTCCATCT CAAGGCATTG TGTGTTTTGT TCCGGGACTG GTTTGGCTGG	3066
GACAAAGTTA GAACTGCCTG AAGTTCGCAC ATTCAGATTG TTGTGTCCAT GGAGTTTTAG	3126
GAGGGGATGG CCTTTCCGGT CTTGCACTT CCATCCTCTC CCCACTTCCC ATCTGGCGTC	3186
CCACACCTTG TCCCCCTGCA CTTCTGGATG ACCAGGGTGC TGCTGCCTCC TAGTCTTTGC	3246
CTTTGCTGGG CTTTCTGTGC AGGAGACTTG GTCTCAAAGC TCAGAGAGAG CCAGTCCGGT	3306
CCCAGCTCCT TTGTCCCTTC CTCAGAGGCC TTCCTTGAAG ATGCATCTAG ACTACCAGCC	3366
TTATCAGTGT TTAAGCTTAT TCCTTTAACA TAAGCTTCCT GACAACATGA AATTGTTGGG	3426

GTTTTGGC GTTGTGAT TTGTTAGGT TTGCTTTAT ACCCGGGCCA AATAGCACAT 3486  
 AACACCTGGT TATATATGAA ATACTCATAT GTTTATGACC AAAATAAATA TGAAACCTCA 3546  
 AAAAAAAAAA AAAAAAAAAA 3566

## (2) INFORMATION FOR SEQ ID NO:2:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 386 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Gly Leu Trp Gly Gln Ser Val Pro Thr Ala Ser Ser Ala Arg Ala  
 -55 -50 -45 -40  
 Gly Arg Tyr Pro Gly Ala Arg Thr Ala Ser Gly Thr Arg Pro Trp Leu  
 -35 -30 -25  
 Leu Asp Pro Lys Ile Leu Lys Phe Val Val Phe Ile Val Ala Val Leu  
 -20 -15 -10  
 Leu Pro Val Arg Val Asp Ser Ala Thr Ile Pro Arg Gln Asp Glu Val  
 -5 1 5  
 Pro Gln Gln Thr Val Ala Pro Gln Gln Gln Arg Arg Ser Leu Lys Glu  
 10 15 20 25  
 Glu Glu Cys Pro Ala Gly Ser His Arg Ser Glu Tyr Thr Gly Ala Cys  
 30 35 40  
 Asn Pro Cys Thr Glu Gly Val Asp Tyr Thr Ile Ala Ser Asn Asn Leu  
 45 50 55  
 Pro Ser Cys Leu Leu Cys Thr Val Cys Lys Ser Gly Gln Thr Asn Lys  
 60 65 70  
 Ser Ser Cys Thr Thr Thr Arg Asp Thr Val Cys Gln Cys Glu Lys Gly  
 75 80 85  
 Ser Phe Gln Asp Lys Asn Ser Pro Glu Met Cys Arg Thr Cys Arg Thr  
 90 95 100 105  
 Gly Cys Pro Arg Gly Met Val Lys Val Ser Asn Cys Thr Pro Arg Ser  
 110 115 120  
 Asp Ile Lys Cys Lys Asn Glu Ser Ala Ala Ser Ser Thr Gly Lys Thr  
 125 130 135  
 Pro Ala Ala Glu Glu Thr Val Thr Thr Ile Leu Gly Met Leu Ala Ser  
 140 145 150  
 Pro Tyr His Tyr Leu Ile Ile Ile Val Val Leu Val Ile Ile Leu Ala  
 155 160 165

Val Val Val Val Gly Phe Ser Cys Arg Lys Lys Phe Ile Ser Tyr Leu  
 170 175 180 185  
 Lys Gly Ile Cys Ser Gly Gly Gly Gly Gly Pro Glu Arg Val His Arg  
 190 195 200  
 Val Leu Phe Arg Arg Arg Ser Cys Pro Ser Arg Val Pro Gly Ala Glu  
 205 210 215  
 Asp Asn Ala Arg Asn Glu Thr Leu Ser Asn Arg Tyr Leu Gln Pro Thr  
 220 225 230  
 Gln Val Ser Glu Gln Glu Ile Gln Gly Gln Glu Leu Ala Glu Leu Thr  
 235 240 245  
 Gly Val Thr Val Glu Ser Pro Glu Glu Pro Gln Arg Leu Leu Glu Gln  
 250 255 260 265  
 Ala Glu Ala Glu Gly Cys Gln Arg Arg Arg Leu Leu Val Pro Val Asn  
 270 275 280  
 Asp Ala Asp Ser Ala Asp Ile Ser Thr Leu Leu Asp Ala Ser Ala Thr  
 285 290 295  
 Leu Glu Glu Gly His Ala Lys Glu Thr Ile Gln Asp Gln Leu Val Gly  
 300 305 310  
 Ser Glu Lys Leu Phe Tyr Glu Glu Asp Glu Ala Gly Ser Ala Thr Ser  
 315 320 325  
 Cys Leu  
 330

## (2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 331 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Met Leu Gly Ile Trp Thr Leu Leu Pro Leu Val Leu Thr Ser Val Ala  
 1 5 10 15  
 Arg Leu Ser Ser Lys Ser Val Asn Ala Gln Val Thr Asp Ile Asn Ser  
 20 25 30  
 Lys Gly Leu Glu Leu Arg Lys Thr Val Thr Val Glu Thr Gln Asn Leu  
 35 40 45  
 Glu Gly Leu His His Asp Gly Gln Phe Cys His Pro Cys Pro Pro Gly  
 50 55 60



Glu Arg Lys Ala Arg Asp Cys Thr Val Asn Gly Asp Glu Pro Asp Cys  
 65 70 75 80  
 Val Pro Cys Gln Glu Gly Lys Glu Tyr Thr Asp Lys Ala His Phe Ser  
 85 90 95  
 Ser Lys Cys Arg Arg Cys Arg Leu Cys Asp Glu Gly His Gly Leu Glu  
 100 105 110  
 Val Glu Ile Asn Cys Thr Arg Thr Gln Asn Thr Lys Cys Arg Cys Lys  
 115 120 125  
 Pro Asn Phe Phe Cys Asn Ser Thr Val Cys Glu His Cys Asp Pro Cys  
 130 135 140  
 Thr Lys Cys Glu His Gly Ile Ile Lys Glu Cys Thr Leu Thr Ser Asn  
 145 150 155 160  
 Thr Lys Cys Lys Glu Glu Gly Ser Arg Ser Asn Gly Trp Leu Cys Leu  
 165 170 175  
 Leu Leu Leu Pro Ile Pro Leu Ile Val Trp Val Lys Arg Lys Glu Val  
 180 185 190  
 Gln Lys Thr Cys Arg Lys His Arg Lys Glu Asn Gln Gly Ser His Glu  
 195 200 205  
 Ser Pro Thr Leu Asn Pro Glu Thr Val Ala Ile Asn Leu Ser Asp Val  
 210 215 220  
 Asp Leu Ser Lys Tyr Ile Thr Thr Ile Ala Gly Val Met Thr Leu Ser  
 225 230 235 240  
 Gln Val Lys Gly Phe Val Arg Lys Asn Gly Val Asn Glu Ala Lys Ile  
 245 250 255  
 Asp Glu Ile Lys Asn Asp Asn Val Gln Asp Thr Ala Glu Gln Lys Val  
 260 265 270  
 Gln Leu Leu Arg Asn Trp His Gln Leu His Gly Lys Lys Glu Ala Tyr  
 275 280 285  
 Asp Thr Leu Ile Lys Asp Leu Lys Lys Ala Asn Leu Cys Thr Leu Ala  
 290 295 300  
 Glu Lys Ile Thr Ile Ile Leu Lys Asp Ile Thr Ser Asp Ser Glu Asn  
 305 310 315 320  
 Ser Asn Phe Arg Asn Glu Ile Gln Ser Leu Val  
 325 330

## (2) INFORMATION FOR SEQ ID NO:4:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 427 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Met	Gly	Ala	Gly	Ala	Thr	Gly	Arg	Ala	Met	Asp	Gly	Pro	Arg	Leu	Leu	1	5	10	15
Leu	Leu	Leu	Leu	Leu	Gly	Val	Ser	Leu	Gly	Gly	Ala	Lys	Glu	Ala	Cys	20	25	30	
Pro	Thr	Gly	Leu	Tyr	Thr	His	Ser	Gly	Glu	Cys	Cys	Lys	Ala	Cys	Asn	35	40	45	
Leu	Gly	Glu	Gly	Val	Ala	Gln	Pro	Cys	Gly	Ala	Asn	Gln	Thr	Val	Cys	50	55	60	
Glu	Pro	Cys	Leu	Asp	Ser	Val	Thr	Phe	Ser	Asp	Val	Val	Ser	Ala	Thr	65	70	75	80
Glu	Pro	Cys	Lys	Pro	Cys	Thr	Glu	Cys	Val	Gly	Leu	Gln	Ser	Met	Ser	85	90	95	
Ala	Pro	Cys	Val	Glu	Ala	Asp	Asp	Ala	Val	Cys	Arg	Cys	Ala	Tyr	Gly	100	105	110	
Tyr	Tyr	Gln	Asp	Glu	Thr	Thr	Gly	Arg	Cys	Glu	Ala	Cys	Arg	Val	Cys	115	120	125	
Glu	Ala	Gly	Ser	Gly	Leu	Val	Phe	Ser	Cys	Gln	Asp	Lys	Gln	Asn	Thr	130	135	140	
Val	Cys	Glu	Glu	Cys	Pro	Asp	Gly	Thr	Tyr	Ser	Asp	Glu	Ala	Asn	His	145	150	155	160
Val	Asp	Pro	Cys	Leu	Pro	Cys	Thr	Val	Cys	Glu	Asp	Thr	Glu	Arg	Gln	165	170	175	
Leu	Arg	Glu	Cys	Thr	Arg	Trp	Ala	Asp	Ala	Glu	Cys	Glu	Glu	Ile	Pro	180	185	190	
Gly	Arg	Trp	Ile	Thr	Arg	Ser	Thr	Pro	Pro	Glu	Gly	Ser	Asp	Ser	Thr	195	200	205	
Ala	Pro	Ser	Thr	Gln	Glu	Pro	Glu	Ala	Pro	Pro	Glu	Gln	Asp	Leu	Ile	210	215	220	
Ala	Ser	Thr	Val	Ala	Gly	Val	Val	Thr	Thr	Val	Met	Gly	Ser	Ser	Gln	225	230	235	240
Pro	Val	Val	Thr	Arg	Gly	Thr	Thr	Asp	Asn	Leu	Ile	Pro	Val	Tyr	Cys	245	250	255	
Ser	Ile	Leu	Ala	Ala	Val	Val	Val	Gly	Leu	Val	Ala	Tyr	Ile	Ala	Phe	260	265	270	
Lys	Arg	Trp	Asn	Ser	Cys	Lys	Gln	Asn	Lys	Gln	Gly	Ala	Asn	Ser	Arg	275	280	285	

Pro Val Asn Gln Thr Pro Pro Pro Glu Gly Glu Lys Leu His Ser Asp  
 290 295 300  
 Ser Gly Ile Ser Val Asp Ser Gln Ser Leu His Asp Gln Gln Pro His  
 305 310 315 320  
 Thr Gln Thr Ala Ser Gly Gln Ala Leu Lys Gly Asp Gly Gly Leu Tyr  
 325 330 335  
 Ser Ser Leu Pro Pro Ala Lys Arg Glu Glu Val Glu Lys Leu Leu Asn  
 340 345 350  
 Gly Ser Ala Gly Asp Thr Trp Arg His Leu Ala Gly Glu Leu Gly Tyr  
 355 360 365  
 Gln Pro Glu His Ile Asp Ser Phe Thr His Glu Ala Cys Pro Val Arg  
 370 375 380  
 Ala Leu Leu Ala Ser Trp Ala Thr Gln Asp Ser Ala Thr Leu Asp Ala  
 385 390 395 400  
 Leu Leu Ala Ala Leu Arg Arg Ile Gln Arg Ala Asp Leu Val Glu Ser  
 405 410 415  
 Leu Cys Ser Glu Ser Thr Ala Thr Ser Pro Val  
 420 425

## (2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 453 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Met Gly Leu Ser Thr Val Pro Asp Leu Leu Leu Pro Leu Val Leu Glu  
 1 5 10 15  
 Leu Leu Val Gly Ile Tyr Pro Ser Gly Val Ile Gly Leu Val Pro His  
 20 25 30  
 Leu Gly Asp Arg Glu Lys Arg Asp Ser Val Cys Pro Gln Gly Lys Tyr  
 35 40 45  
 Ile His Pro Asn Asn Ser Ile Cys Cys Thr Lys Cys His Lys Gly Thr  
 50 55 60  
 Tyr Leu Tyr Asn Asp Cys Pro Gly Pro Gly Gln Asp Thr Asp Cys Arg  
 65 70 75 80  
 Glu Cys Glu Ser Gly Ser Phe Thr Ala Ser Glu Asn His Leu Arg His  
 85 90 95

Cys Leu Ser Cys Ser Lys Cys Arg Lys Glu Met Gly Gln Val Glu Ile  
 100 105 110  
 Ser Ser Cys Thr Val Asp Arg Asp Thr Val Cys Gly Cys Arg Lys Asn  
 115 120 125  
 Gln Tyr Arg His Tyr Trp Ser Glu Asn Leu Phe Gln Cys Phe Asn Cys  
 130 135 140  
 Ser Leu Cys Leu Asn Gly Thr Val His Leu Ser Cys Gln Glu Lys Gln  
 145 150 155 160  
 Asn Thr Val Cys Thr Cys His Ala Gly Phe Phe Leu Arg Glu Asn Glu  
 165 170 175  
 Cys Val Ser Cys Ser Asn Cys Lys Lys Ser Leu Glu Cys Thr Lys Leu  
 180 185 190  
 Cys Leu Pro Gln Ile Glu Asn Val Lys Gly Thr Glu Asp Ser Gly Thr  
 195 200 205  
 Thr Val Leu Leu Pro Leu Val Ile Phe Phe Gly Leu Cys Leu Leu Ser  
 210 215 220  
 Leu Leu Phe Ile Gly Leu Met Tyr Arg Tyr Gln Arg Trp Lys Ser Lys  
 225 230 235 240  
 Leu Tyr Ser Ile Val Cys Gly Lys Ser Thr Pro Glu Lys Glu Gly Glu  
 245 250 255  
 Leu Glu Gly Thr Thr Thr Lys Pro Leu Ala Pro Asn Pro Ser Phe Ser  
 260 265 270  
 Pro Thr Pro Gly Phe Thr Pro Thr Leu Gly Phe Ser Pro Val Pro Ser  
 275 280 285  
 Ser Thr Phe Thr Ser Ser Ser Thr Tyr Thr Pro Gly Asp Cys Pro Asn  
 290 295 300  
 Phe Ala Ala Pro Arg Arg Glu Val Ala Pro Pro Tyr Gln Gly Ala Asp  
 305 310 315 320  
 Pro Ile Leu Ala Thr Ala Leu Ala Ser Asp Pro Ile Pro Asn Pro Leu  
 325 330 335  
 Gln Lys Trp Glu Asp Ser Ala His Lys Pro Gln Ser Leu Asp Thr Asp  
 340 345 350  
 Asp Pro Ala Thr Leu Tyr Ala Val Val Glu Asn Val Pro Pro Leu Arg  
 355 360 365  
 Trp Lys Glu Phe Val Arg Arg Leu Gly Leu Ser Asp His Glu Ile Asp  
 370 375 380  
 Arg Leu Glu Leu Gln Asn Gly Arg Cys Leu Arg Glu Ala Gln Tyr Ser  
 385 390 395 400  
 Met Leu Ala Thr Trp Arg Arg Arg Thr Pro Arg Arg Glu Ala Thr Leu  
 405 410 415

Glu Leu Leu Gly Arg Val Leu Arg Asp Met Asp Leu Leu Gly Cys Leu  
 420 425 430

Glu Asp Ile Glu Glu Ala Leu Cys Gly Pro Ala Ala Leu Pro Pro Ala  
 435 440 445

Pro Ser Leu Leu Arg  
 450

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 467 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Met Ala Pro Pro Pro Ala Arg Val His Leu Gly Ala Phe Leu Ala Val  
 1 5 10 15

Thr Pro Asn Pro Gly Ser Ala Ala Ser Gly Thr Glu Ala Ala Ala Ala  
 20 25 30

Thr Pro Ser Lys Val Trp Gly Ser Ser Ala Gly Arg Ile Glu Pro Arg  
 35 40 45

Gly Gly Gly Arg Gly Ala Leu Pro Thr Ser Met Gly Gln His Gly Pro  
 50 55 60

Ser Ala Arg Ala Arg Ala Gly Arg Ala Pro Gly Pro Arg Pro Ala Arg  
 65 70 75 80

Glu Ala Ser Pro Arg Leu Arg Val His Lys Thr Phe Lys Phe Val Val  
 85 90 95

Val Gly Val Leu Leu Gln Val Val Pro Ser Ser Ala Ala Thr Ile Lys  
 100 105 110

Leu His Asp Gln Ser Ile Gly Thr Gln Gln Trp Glu His Ser Pro Leu  
 115 120 125

Gly Glu Leu Cys Pro Pro Gly Ser His Arg Ser Glu Arg Pro Gly Ala  
 130 135 140

Cys Asn Arg Cys Thr Glu Gly Val Gly Tyr Thr Asn Ala Ser Asn Asn  
 145 150 155 160

Leu Phe Ala Cys Leu Pro Cys Thr Ala Cys Lys Ser Asp Glu Glu Glu  
 165 170 175

Arg Ser Pro Cys Thr Thr Thr Arg Asn Thr Ala Cys Gln Cys Lys Pro  
 180 185 190

Gly Thr Phe Arg Asn Asp Asn Ser Ala Glu Met Cys Arg Lys Cys Ser  
 195 200 205  
 Thr Gly Cys Pro Arg Gly Met Val Lys Val Lys Asp Cys Thr Pro Trp  
 210 215 220  
 Ser Asp Ile Glu Cys Val His Lys Glu Ser Gly Asn Gly His Asn Ile  
 225 230 235 240  
 Trp Val Ile Leu Val Val Thr Leu Val Val Pro Leu Leu Leu Val Ala  
 245 250 255  
 Val Leu Ile Val Cys Cys Cys Ile Gly Ser Gly Cys Gly Gly Asp Pro  
 260 265 270  
 Lys Cys Met Asp Arg Val Cys Phe Trp Arg Leu Gly Leu Leu Arg Gly  
 275 280 285  
 Pro Gly Ala Glu Asp Asn Ala His Asn Glu Ile Leu Ser Asn Ala Asp  
 290 295 300  
 Ser Leu Ser Thr Phe Val Ser Glu Gln Gln Met Glu Ser Gln Glu Pro  
 305 310 315 320  
 Ala Asp Leu Thr Gly Val Val Gln Ser Pro Gly Glu Ala Gln Cys Leu  
 325 330 335  
 Leu Gly Pro Ala Glu Ala Glu Gly Ser Gln Arg Arg Arg Leu Leu Val  
 340 345 350  
 Pro Ala Asn Gly Ala Asp Pro Thr Glu Thr Leu Met Leu Phe Phe Asp  
 355 360 365  
 Lys Phe Ala Asn Ile Val Pro Phe Asp Ser Trp Asp Gln Leu Met Arg  
 370 375 380  
 Gln Leu Asp Leu Thr Lys Asn Glu Ile Asp Val Val Arg Ala Gly Thr  
 385 390 395 400  
 Ala Gly Pro Gly Asp Ala Leu Tyr Ala Met Leu Met Lys Trp Val Asn  
 405 410 415  
 Lys Thr Gly Arg Asn Ala Ser Ile His Thr Leu Leu Asp Ala Leu Glu  
 420 425 430  
 Arg Met Glu Glu Arg His Ala Lys Glu Lys Ile Gln Asp Leu Leu Val  
 435 440 445  
 Asp Ser Gly Lys Phe Ile Tyr Leu Glu Asp Gly Thr Gly Ser Ala Val  
 450 455 460  
 Ser Leu Glu  
 465

## (2) INFORMATION FOR SEQ ID NO:7:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 343 base pairs

(B) TYPE: nucleic acid



- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

GTCACGTTCC ATTATTTTAT TTAAAAACCT CAGTAATCGT TTTAGCTTCT TTCCAGCAAA	60
CTCTTCTCCA CAGTAGCCCA GTCGTGGTAG GATAAATTAC GGATATAGTC ATTCTAGGGG	120
TTTCAGTCTT TTCCATCTCA AGGCATTGTG TGTTTTGTTT CGGGACTGGT TTGGCTGGGA	180
CAAAGTTAGA ACTGCCTGAA GTTCGCACAT TCAGATTGTT GTGTCCATGG AGTTTTAGGA	240
GGGGATGGCC TTTCCGGTCT TCGCACTTCC ATCCTCTCCC ACTTCCATCT GGCCTCCACA	300
ACTTGTCCCC TGCACTTCTG GATGACACAG GGTGCTGCTG CCT	343

(2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 279 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

GTGGACGGTG CTTGGGAATG GTGTGAAATG GTTGCCGACT CAGGCATGGA TGGGCCCCCTC	60
TCGCTTCTGG TGGTCTGTGA ACTGAGTCCC TGGGATGCCT TTAGGGCAGA GATTCCTGAG	120
CTGCGTTTTA GGTACAGAT TCCCTGTTTG AGGAGCTTGG CCCCTCTGTA AGCGTCTGAC	180
TCATCTCAGA GATATCAATT CTTAAACACT GTGACAACGG GATCTAAAAT GGCTGACACA	240
TTTGTCTTGG TGTCACGTTT CATTATTTTA TTTAAAATT	279

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 250 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

GGCCACGTAG TGCCACGTGC CACAACTAC GGGGGACGAT TTCTGATTGA ATTTTGGCG 60  
CTTTCATCC ACCCTCCTCC CTTCTAATGG GACTTTGGGG ACAAGGTCC GACCGCCTCG 120  
AGCGTCGACA GGGCGCTATC CAGGAGCCAG GACAGCGTCG GGAACCAGAC CATGGCTCCT 180  
GGACCCCAAG ATCCTTAAGT TCGTCGTCTT CATCGTCGGG TTCTCTGCCG GTAAGTTAGG 240  
AGGTCCCTGG 250

## (2) INFORMATION FOR SEQ ID NO:10:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

CGCCCATGGC CACCATCCCC CGGCAG 26

## (2) INFORMATION FOR SEQ ID NO:11:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

CGCAAGCTTT TAGTAGTGAT AGGGAGAGGC 30

## (2) INFORMATION FOR SEQ ID NO:12:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 33 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

CGCGGATCCG CCATCATGGG ACTTTGGGGA CAA

33

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

CGCGGTACCT TAGTAGTGAT AGGGAGAGGC

30

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 58 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

CGCTCTAGAT CAAGCGTAGT CTGGGACGTC GTATGGGTAG TAAGTGATAG GGAGAGGC

58

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 498 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)


(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

GAGTTTGACC AGAGATGCAA GGGGTGAAGG AGCGCTTCCT ACCGTTAGGA ACTCTGGGGA	60
CAGAGCGCCC CGGCCGCCTG ATGGCGAGGC AGGGTGCGAC CCAGGACCCA GGACGGCGTC	120
GGGAACCATA CCATGGCCCG GATCCCCAAG ACCCTAAAGT TCGTCGTCGT CATCGTCGCG	180
GTCTTGCTGC CAGTCCTAGC TTA CTCTGCC ACCACTGCCC GGCAGAGGGA AGTTCCCCAG	240
CAGACAGTGG CCCCACAGCA ACAGAGGCAC AGCTTCAAGG GGGAGGAGTG TCCAGCAGGA	300
TCTCATAGAT CAGAACATAC TGGAGCCTGT AACCCGTGCA CAGAGGGTGT GGATTACACC	360
AACGCTTCCA ACAATGAACC TTCTTGCTTC CCATGTAC	498

## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

<b>A.</b> The indications made below relate to the microorganism referred to in the description on page <u>4</u> , line <u>15</u>	
<b>B. IDENTIFICATION OF DEPOSIT</b> <span style="float: right;">Further deposits are identified on an additional sheet <input type="checkbox"/></span>	
Name of depositary institution <div style="text-align: center;">American Type Culture Collection</div>	
Address of depositary institution ( <i>including postal code and country</i> ) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit    May 15, 1997	Accession Number    209040
<b>C. ADDITIONAL INDICATIONS</b> ( <i>leave blank if not applicable</i> ) <span style="float: right;">This information is continued on an additional sheet <input type="checkbox"/></span>	
<b>D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE</b> ( <i>If the indications are not for all designated States</i> )	
<b>E. SEPARATE FURNISHING OF INDICATIONS</b> ( <i>leave blank if not applicable</i> )	
The indications listed below will be submitted to the International Bureau later ( <i>specify the general nature of the indications, e.g., "Accession Number of Deposit"</i> )	

<div style="border-bottom: 1px solid black; margin-bottom: 5px;">For receiving Office use only</div> <div style="display: flex; align-items: center; margin-bottom: 10px;"><input checked="" type="checkbox"/> This sheet was received with the international application</div> <div style="border-bottom: 1px solid black; margin-bottom: 10px;">Authorized officer</div> <div style="text-align: center;"></div>	<div style="border-bottom: 1px solid black; margin-bottom: 5px;">For International Bureau use only</div> <div style="display: flex; align-items: center; margin-bottom: 10px;"><input type="checkbox"/> This sheet was received by the International Bureau on:</div> <div style="border-bottom: 1px solid black; margin-bottom: 10px;">Authorized officer</div>
---	--

*What Is Claimed Is:*

1. An isolated nucleic acid molecule comprising a polynucleotide having a nucleotide sequence at least 95% identical to a sequence selected from the group consisting of:

- (a) a nucleotide sequence encoding a polypeptide comprising amino acids from about -55 to about 331 in SEQ ID NO:2;
  - (b) a nucleotide sequence encoding a polypeptide comprising amino acids from about -54 to about 331 in SEQ ID NO:2;
  - (c) a nucleotide sequence encoding a polypeptide comprising amino acids from about 1 to about 331 in SEQ ID NO:2;
  - (d) a nucleotide sequence encoding a polypeptide having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. 209040;
  - (e) a nucleotide sequence encoding the mature TR10 polypeptide having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. 209040;
  - (f) a nucleotide sequence encoding the TR10 extracellular domain;
  - (g) a nucleotide sequence encoding the TR10 transmembrane domain;
  - (h) a nucleotide sequence encoding the TR10 intracellular domain;
  - (i) a nucleotide sequence encoding the TR10 receptor extracellular and intracellular domains with all or part of the transmembrane domain deleted;
  - (j) a nucleotide sequence encoding the partial TR10 death domain;
- and
- (k) a nucleotide sequence complementary to any of the nucleotide sequences in (a), (b), (c), (d), (e), (f), (g), (h), (i), or (j).

2. The nucleic acid molecule of claim 1, wherein said polynucleotide has the nucleotide sequence in SEQ ID NO:1.

3. The nucleic acid molecule of claim 1, wherein said polynucleotide has the nucleotide sequence in SEQ ID NO:1 encoding the TR10 receptor having the amino acid sequence in SEQ ID NO:2.

4. The nucleic acid molecule of claim 1, wherein said polynucleotide has the nucleotide sequence in SEQ ID NO:1 encoding the mature TR10 receptor having the amino acid sequence in SEQ ID NO:2.

5. The nucleic acid molecule of claim 1, wherein said polynucleotide has the complete nucleotide sequence of the cDNA clone contained in ATCC Deposit No. 209040.

6. The nucleic acid molecule of claim 1, wherein said polynucleotide has the nucleotide sequence encoding the TR10 receptor having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. 209040.

7. The nucleic acid molecule of claim 1, wherein said polynucleotide has the nucleotide sequence encoding the mature TR10 receptor having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. 209040.

8. An isolated nucleic acid molecule comprising a polynucleotide which hybridizes under stringent hybridization conditions to a polynucleotide having a nucleotide sequence identical to a nucleotide sequence in (a), (b), (c), (d), (e), (f), (g), (h), (i), (j), or (k) of claim 1, wherein said polynucleotide which hybridizes does not hybridize under stringent hybridization conditions to a polynucleotide having a nucleotide sequence consisting of only A residues or of only T residues.

9. An isolated nucleic acid molecule comprising a polynucleotide which encodes the amino acid sequence of an epitope-bearing portion of a TR10 receptor having an amino acid sequence in (a), (b), (c), (d), (e), (f), (g), (h), (i), or (j) of claim 1.

10. The isolated nucleic acid molecule of claim 9, which encodes an epitope-bearing portion of a TR10 receptor selected from the group consisting of: a polypeptide comprising amino acid residues from about 2 to about 58 in SEQ ID NO:2; a polypeptide comprising amino acid residues from about 75 to about 142 in SEQ ID NO:2; a polypeptide comprising amino acid residues from about 195 to about 228 in SEQ ID NO:2.



11. The isolated nucleic acid molecule of claim 1, which encodes the TR10 receptor extracellular domain.

12. The isolated nucleic acid molecule of claim 1, which encodes the TR10 receptor transmembrane domain.

13. The isolated nucleic acid molecule of claim 1, which encodes the TR10 receptor intracellular domain.

14. An isolated nucleic acid molecule comprising a polynucleotide having a sequence at least 95% identical to a sequence selected from the group consisting of:

- (a) the nucleotide sequence of cloneHSABD50R (SEQ ID NO:7);
- (b) the nucleotide sequence of cloneHGBDL20R (SEQ ID NO:8);
- (c) the nucleotide sequence of cloneHELDL61R (SEQ ID NO:9);

and

(d) a nucleotide sequence complementary to any of the nucleotide sequences in (a), (b), or (c) above.

15. A method for making a recombinant vector comprising inserting an isolated nucleic acid molecule of claim 1 into a vector.

16. A recombinant vector produced by the method of claim 15.

17. A method of making a recombinant host cell comprising introducing the recombinant vector of claim 16 into a host cell.

18. A recombinant host cell produced by the method of claim 17.

19. A recombinant method for producing a TR10 polypeptide, comprising culturing the recombinant host cell of claim 18 under conditions such that said polypeptide is expressed, and recovering said polypeptide.

20. An isolated TR10 polypeptide having an amino acid sequence at least 95% identical to a sequence selected from the group consisting of:

- (a) amino acids from about -55 to about 331 in SEQ ID NO:2;
- (b) amino acids from about -54 to about 331 in SEQ ID NO:2;

- (c) amino acids from about 1 to about 331 in SEQ ID NO:2;
- (d) the amino acid sequence of the TR10 polypeptide having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. 209040;
- (e) the amino acid sequence of the mature TR10 polypeptide having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. 209040;
- (f) the amino acid sequence of the TR10 receptor extracellular domain;
- (g) the amino acid sequence of the TR10 receptor transmembrane domain;
- (h) the amino acid sequence of the TR10 receptor intracellular domain;
- (i) the amino acid sequence of the TR10 receptor intracellular and extracellular domains with all or part of the transmembrane domain deleted;
- (j) the amino acid sequence of the TR10 receptor death domain; and
- (k) the amino acid sequence of an epitope-bearing portion of any one of the polypeptides of (a), (b), (c), (d), (e), (f), (g), (h), (i), or (j).

21. An isolated polypeptide comprising an epitope-bearing portion of the TR10 receptor protein, wherein said portion is selected from the group consisting of: a polypeptide comprising amino acid residues from about 2 to about 58 in SEQ ID NO:2; a polypeptide comprising amino acid residues from about 75 to about 142 in SEQ ID NO:2; and a polypeptide comprising amino acid residues from about 195 to about 228 in SEQ ID NO:2

22. An isolated antibody that binds specifically to a TR10 receptor polypeptide of claim 20.

23. A method of treating diseases and disorders associated with the inhibition of apoptosis comprising administering an effective amount of the polypeptide as claimed in claim 20, or an agonist thereof to a patient in need thereof.

24. A method of treating diseases and disorders associated with increased apoptosis comprising administering to a patient in need thereof an effective amount of an antagonist of the polypeptide as claimed in claim 20 to a patient in need thereof.

25. A method of treating inflammatory diseases and disorders comprising administering to a patient in need thereof an effective amount of an antagonist of the polypeptide as claimed in claim 20.

26. An isolated nucleic acid molecule comprising a polynucleotide encoding a TR10 receptor polypeptide wherein, except for at least one conservative amino acid substitution, said polypeptide has a sequence selected from the group consisting of:

- (a) a nucleotide sequence encoding a polypeptide comprising amino acids from about -55 to about 331 in SEQ ID NO:2;
  - (b) a nucleotide sequence encoding a polypeptide comprising amino acids from about -54 to about 331 in SEQ ID NO:2;
  - (c) a nucleotide sequence encoding a polypeptide comprising amino acids from about 1 to about 331 in SEQ ID NO:2;
  - (d) a nucleotide sequence encoding a polypeptide having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. 209040;
  - (e) a nucleotide sequence encoding the mature TR10 polypeptide having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. 209040;
  - (f) a nucleotide sequence encoding the TR10 extracellular domain;
  - (g) a nucleotide sequence encoding the TR10 transmembrane domain;
  - (h) a nucleotide sequence encoding the TR10 intracellular domain;
  - (i) a nucleotide sequence encoding the TR10 receptor extracellular and intracellular domains with all or part of the transmembrane domain deleted;
  - (j) a nucleotide sequence encoding the TR10 partial death domain;
- and
- (k) a nucleotide sequence complementary to any of the nucleotide sequences in (a), (b), (c), (d), (e), (f), (g), (h), (i), or (j).

27. An isolated TR10 receptor polypeptide wherein, except for at least one conservative amino acid substitution, said polypeptide has a sequence selected from the group consisting of:

- (a) amino acids from about -55 to about 331 in SEQ ID NO:2;
- (b) amino acids from about -54 to about 331 in SEQ ID NO:2;
- (c) amino acids from about 1 to about 331 in SEQ ID NO:2;

- (d) the amino acid sequence of the TR10 polypeptide having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. 209040;
- (e) the amino acid sequence of the mature TR10 polypeptide having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. 209040;
- (f) the amino acid sequence of the TR10 receptor extracellular domain;
- (g) the amino acid sequence of the TR10 receptor transmembrane domain;
- (h) the amino acid sequence of the TR10 receptor intracellular domain;
- (i) the amino acid sequence of the TR10 receptor extracellular and intracellular domains with all or part of the transmembrane domain deleted;
- (j) the amino acid sequence of the TR10 receptor partial death domain; and
- (k) the amino acid sequence of an epitope-bearing portion of any one of the polypeptides of (a), (b), (c), (d), (e), (f), (g), (h), (i), or (j).

1/7

10 30 50  
CGACCCACGCGTCCGCCCACGCGTCCGGAGAACCTTTGCACGCGCACAACTACGGGGAC  
70 90 110  
GATTTCTGATTGATTTTGGCGCTTTCGATCCACCCTCCTCCCTTCTCATGGGACTTTGG  
M G L W  
130 150 170  
GGACAAAGCGTCCCGACCGCCTCGAGCGCTCGAGCAGGGCGCTATCCAGGAGCCAGGACA  
G Q S V P T A S S A R A G R Y P G A R T  
190 210 230  
GCGTCGGGAACCAGACCATGGCTCCTGGACCCCAAGATCCTTAAGTTCGTCGTCTTCATC  
A S G T R P W L L D P K I L K F V V F I  
250 270 290  
GTCGCGGTTCTGCTGCCGGTCCGGGTGACTCTGCCACCATCCCCGGCAGGACGAAGTT  
V A V L L P V R V D S A T I P R Q D E V  
310 330 350  
CCCCAGCAGACAGTGGCCCCACAGCAACAGAGGGCGCAGCCTCAAGGAGGAGGAGTGTCCA  
P Q Q T V A P Q Q Q R R S L K E E E C P  
370 390 410  
GCAGGATCTCATAGATCAGAATATACTGGAGCCTGTAACCCGTGCACAGAGGGTGTGGAT  
A G S H R S E Y T G A C N P C T E G V D  
430 450 470  
TACACCATTGCTTCCAACAATTTGCCTTCTTGCTGCTATGTACAGTTTGTAAATCAGGT  
Y T I A S N N L P S C L L C T V C K S G  
490 510 530  
CAAACAAATAAAAGTTCCTGTACCACGACCAGAGACACCGTGTGTGAGTGTGAAAAAGGA  
Q T N K S S C T T T R D T V C Q C E K G  
550 570 590  
AGCTTCCAGGATAAAAACTCCCCTGAGATGTGCCGGACGTGTAGAACAGGGTGTCCCAGA  
S F Q D K N S P E M C R T C R T G C P R  
610 630 650  
GGGATGGTCAAGGTCAGTAATTGTACGCCCCGGAGTGACATCAAGTGCAAAAATGAATCA  
G M V K V S N C T P R S D I K C K N E S  
670 690 710  
GCTGCCAGTTCCACTGGGAAAACCCAGCAGCGGAGGAGACAGTGACCACCATCCTGGGG  
A A S S T G K T P A A E E T V T T I L G  
730 750 770  
ATGCTTGCCTCTCCCTATCACTACCTTATCATCATAGTGGTTTTAGTCATCATTTTAGCT  
M L A S P Y H Y L I I I V V L V I I L A

FIG.1A

SUBSTITUTE SHEET (RULE 26)

2/7

790 810 830  
GTGGTTGTGGTTGGCTTTTCATGTCGGAAGAAATTCATTTCTTACCTCAAAGGCATCTGC  
V V V V G F S C R K K F I S Y L K G I C  
850 870 890  
TCAGGTGGTGGAGGAGGTCCCGAACGTGTGCACAGAGTCCTTTTCCGGCGGCGTTCATGT  
S G G G G G P E R V H R V L F R R R S C  
910 930 950  
CCTTCACGAGTTCCTGGGGCGGAGGACAATGCCCGCAACGAGACCCTGAGTAACAGATAC  
P S R V P G A E D N A R N E T L S N R Y  
970 990 1010  
TTGCAGCCCACCCAGGTCTCTGAGCAGGAAATCCAAGGTCAGGAGCTGGCAGAGCTAACA  
L Q P T Q V S E Q E I Q G Q E L A E L T  
1030 1050 1070  
GGTGTGACTGTAGAGTCGCCAGAGGAGCCACAGCGTCTGCTGGAACAGGCAGAAGCTGAA  
G V T V E S P E E P Q R L L E Q A E A E  
1090 1110 1130  
GGGTGTCAGAGGAGGAGGCTGCTGGTTCCAGTGAATGACGCTGACTCCGCTGACATCAGC  
G C Q R R R L L V P V N D A D S A D I S  
1150 1170 1190  
ACCTTGCTGGATGCCTCGGCAACACTGGAAGAAGGACATGCAAAGGAAACAATTCAGGAC  
T L L D A S A T L E E G H A K E T I Q D  
1210 1230 1250  
CAACTGGTGGGCTCCGAAAAGCTCTTTTATGAAGAAGATGAGGCAGGCTCTGCTACGTCC  
Q L V G S E K L F Y E E D E A G S A T S  
1270 1290 1310  
TGCCTGTGAAAGAATCTCTTCAGGAAACCAGAGCTTCCCTCATTACCTTTTCTCCTACA  
C L \*  
1330 1350 1370  
AAGGGAAGCAGCCTGGAAGAAACAGTCCAGTACTTGACCCATGCCCCAACAACTCTACT  
1390 1410 1430  
ATCCAATATGGGGCAGCTTACCAATGGTCCTAGAACTTTGTTAACGCACTTGAGTAATT  
1450 1470 1490  
TTTATGAAATACTGCGTGTGATAAGCAAACGGGAGAAATTTATATCAGATTCTTGGCTGC  
1510 1530 1550  
ATAGTTATACGATTGTGTATTAAGGGTCGTTTTAGGCCACATGCGGTGGCTCATGCCTGT  
1570 1590 1610  
AATCCCAGCACTTTGATAGGCTGAGGCAGGTGGATTGCTTTGAGCTCGGGAGTTTGAGAC  
1630 1650 1670  
CAGCCTCATCAACACAGTGAAACTCCATCTCAATTTAAAAAGAAAAAAAGTGGTTTTAG

FIG.1B

SUBSTITUTE SHEET (RULE 26)



3/7

1690	1710	1730
GATGTCATTCTTTGCAGTTCTTCATCATGAGACAAGTCTTTTTTCTGCTTCTTATATTG		
1750	1770	1790
CAAGCTCCATCTCTACTGGTGTGTGCATTTAATGACATCTAACTACAGATGCCGCACAGC		
1810	1830	1850
CACAATGCTTTGCCTTATAGTTTTTTAACTTTAGAACGGGATTATCTTGTTATTACCTGT		
1870	1890	1910
ATTTTCAGTTTCCGATATTTTTGACTTAATGATGAGATTATCAAGACGTAGCCCTATGCT		
1930	1950	1970
AAGTCATGAGCATATGGACTTACGAGGGTTCGACTTAGAGTTTTGAGCTTTAAGATAGGA		
1990	2010	2030
TTATTGGGGCTTACCCCCACCTTAATTAGAGAAACATTTATATTGCTTACTACTGTAGGC		
2050	2070	2090
TGTACATCTCTTTTCCGATTTTTGTATAATGATGTAACATGGAAAACTTTAGGAAATG		
2110	2130	2150
CACTTATTAGGCTGTTTACATGGGTTCCTGGATACAAATCAGCAGTCAAAAATGACTAA		
2170	2190	2210
AAATATAACTAGTGACGGAGGGAGAAATCCTCCCTCTGTGGGAGGCACTTACTGCATTCC		
2230	2250	2270
AGTTCTCCCTCCTGCGCCCTGAGACTGGACCAGGGTTTGATGGCTGGCAGCTTCTCAAGG		
2290	2310	2330
GGCAGCTTGTCTTACTTGTTAATTTAGAGGTATATAGCCATATTTATTTATAAATAAAT		
2350	2370	2390
ATTTATTTATTTATTTATAAGTAGATGTTTACATATGCCCAGGATTTTGAAGAGCCTGGT		
2410	2430	2450
ATCTTTGGGAAGCCATGTGTCTGGTTTGTCTGCTGGGACAGTCATGGGACTGCATCTTC		
2470	2490	2510
CGACTTGTCCACAGCAGATGAGGACAGTGAGAATTAAGTTAGATCCGAGACTGCGAAGAG		
2530	2550	2570
CTTCTCTTTCAAGCGCCATTACAGTTGAACGTTAGTGAATCTTGAGCCTCATTGGGCTC		
2590	2610	2630
AGGGCAGAGCAGGTGTTTATCTGCCCCGGCATCTGCCATGGCATCAAGAGGGAAGAGTGG		
2650	2670	2690
ACGGTGCTTGGGAATGGTGTGAAATGGTTGCCGACTCAGGCATGGATGGGCCCCCTCTCGC		
2710	2730	2750
TTCTGGTGGTCTGTGAACTGAGTCCCTGGGATGCCTTTTAGGGCAGAGATTCTGAGCTG		
2770	2790	2810
CGTTTTAGGGTACAGATTCCCTGTTTGAGGAGCTTGGCCCCCTCTGTAAGCATCTGACTCA		

FIG.1C

SUBSTITUTE SHEET (RULE 26)

4/7

2830 2850 2870  
TCTCAGAGATATCAATTCTTAAACACTGTGACAACGGGATCTAAAATGGCTGACACATTT  
2890 2910 2930  
GTCCTTGTGTACGTTCCATTATTTTATTTAAAAACCTCAGTAATCGTTTTAGCTTCTTT  
2950 2970 2990  
CCAGCAAACCTCTTCTCCACAGTAGCCCAGTCGTGGTAGGATAAATTACGGATATAGTCAT  
3010 3030 3050  
TCTAGGGGTTTCAGTCTTTTCCATCTCAAGGCATTGTGTGTTTTGTTCCGGGACTGGTTT  
3070 3090 3110  
GGCTGGGACAAAGTTAGAACTGCCTGAAGTTTCGCACATTCAGATTGTTGTGTCCATGGAG  
3130 3150 3170  
TTTAGGAGGGGATGGCCTTTCCGGTCTTCGCACTTCCATCCTCTCCCCACTTCCCATCT  
3190 3210 3230  
GGCGTCCCACACCTTGTCCTCCCTGCACTTCTGGATGACCAGGGTGCTGCTGCCTCCTAGT  
3250 3270 3290  
CTTTGCCCTTTGCTGGGCCTTCTGTGCAGGAGACTTGGTCTCAAAGCTCAGAGAGAGCCAG  
3310 3330 3350  
TCCGGTCCCAGCTCCTTTGTCCCTTCCTCAGAGGCCTTCCTTGAAGATGCATCTAGACTA  
3370 3390 3410  
CCAGCCTTATCAGTGTTTAAGCTTATTCCTTTAACATAAGCTTCCTGACAACATGAAATT  
3430 3450 3470  
GTTGGGGTTTTTTGGCGTTTGTTGATTGTTTAGGTTTTGCTTTATACCCGGGCCAAATA  
3490 3510 3530  
GCACATAACACCTGGTTATATATGAAATACTCATATGTTTATGACCAAAATAAATATGAA  
3550  
ACCTCAAAAAAAAAAAAAAAAAAAAAA

FIG.1D

**FIG. 2A**

1184 I P L I V - - - W V K R K E V Q K T C R K H R K E N Q G S H E S P T - - - L N - - - P E - - - T V A I - N L S D V D L S K Y I T h F a s  
2231 V V T T V M G S S - - - Q P V V T R G T T D N L I P V Y C S I L A A V V G L - - - V A Y I A F K R W N S C K Q N K Q G A N S R P V N Q T P P h N G F R P  
2230 I G L M Y R Y Q R W - K S K L Y S I V C G K S T P E K E G E L E G T T K P L A - - - P N P S F S P T P G F T P T L - G F S P V P S S T F T S h T N F R I  
2221 I I L A V V - - - - - V V G F S C R K K F I S Y L K G I C S G G G P E R V H R V L F R R R - S C P S F V P G A E D N A R N E T L S T R I O  
2252 L L L V A V - - - - - L I V C C - - - - - I G S C C G G D P K C M D R V C F W R L - G - L L F G P G A E D N A H N E I L S D R - 4

2235T - - - - - I A G V M T L S Q V - - - - - h Fas  
2296P E G E K L H S D S G I S V D S Q S L H D Q Q P H T Q T A S G Q A - - L K G D G G L Y S S - - - - - L P A K R E E V E K L L - - - - - h NGFR P  
2296S S T Y T P G D C P N F A A P R R E V A - - P P Y Q G A D P I L A T A L A S D P I P N P L Q K W E D S A H K P Q S L D T D P A T L Y A V V h TNFR I  
2282N R Y L Q P T Q V S E C E I Q G Q E L A E L T G V T V E S P E E P Q R L L E Q A E A E G C Q R R R L L V P V N D A D S A E - - - - - T R I O  
302N A D S L S T F V S E C Q M E S Q E P A D L T G V T V Q S P G E A C C L L G P A E A E G S Q R R R L L V P A N G A D P T E T L M L F F D K F D R - 4

[illegible]

3306 LA EK IO TI IL KD IT SD SE NS FR NE IQ SL V  
4413 - - - - - LV ES - - - - - LC SE ST AT SP V  
4433 CL E DI EE AL - - - - - CG PA AL PP APS L LR  
4435 AK ET IQ DQ LV GS EK LF YE DE AG SA TS CL  
4440 AK EK IQ DL LV DS GK FI YL ED GT GS AV S - LE

h Fas  
h NGER p  
h TNFR 1  
TR10  
DR-4

**FIG. 2B**

7/7





FIG.3

Applicant's or agent's file reference number	E379PCT	International application
---	---------	---------------------------

## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page 4, line 15	
B. IDENTIFICATION OF DEPOSIT <span style="float: right;">Further deposits are identified on an additional sheet <input type="checkbox"/></span>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit May 15, 1997	Accession Number 209040
C. ADDITIONAL INDICATIONS (leave blank if not applicable) <span style="float: right;">This information is continued on an additional sheet <input type="checkbox"/></span>	
DNA Plasmid 2174510  In respect of those designations in which a European Patent is sought a sample of the deposited microorganism will be available until the publication of the mention of the grant of the European patent or until the date on which the application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28(4)EPC).	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications, e.g., "Accession Number of Deposit")	

For receiving Office use only	For International Bureau use only
<input checked="" type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer 	Authorized officer



## **DNA Plasmid No. 2174510**

### **CANADA**

The applicant requests that, until either a Canadian patent has been issued on the basis of an application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the Commissioner of Patents only authorizes the furnishing of a sample of the deposited biological material referred to in the application to an independent expert nominated by the Commissioner, the applicant must, by a written statement, inform the International Bureau accordingly before completion of technical preparations for publication of the international application.

### **NORWAY**

The applicant hereby requests that the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on the list of recognized experts drawn up by the Norwegian Patent Office or any person approved by the applicant in the individual case.

### **AUSTRALIA**

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

### **FINLAND**

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Regulations), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

### **UNITED KINGDOM**

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for the international publication of the application.

## **DENMARK**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent Office or any person by the applicant in the individual case.

## **SWEDEN**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent Office or any person approved by a applicant in the individual case.

## **NETHERLANDS**

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in the 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US98/10981**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(6) : CO7H 21/04; CO7K 14/705; C12N 15/09, 15/63; C12Q 1/68

US CL : 536/23.5, 24.3; 435/7.2, 69.1, 320.1; 530/350

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 536/23.5, 24.3; 435/7.2, 69.1, 320.1; 530/350

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WALCZAK, H et al. TRAIL-R2: A Novel Apoptosis-Mediating Receptor for TRAIL. The EMBO Journal. 1997, Vol. 16, No. 17, pages 5386-5397.	1-13, 15-21, 26-27
A	MARSTERS, S. A. et al. A Novel receptor for Apo2L/TRAIL Contains a Truncated Death Domain. Current Biology. 1997, Vol 7, No. 12, pages 1003-1006.	1-3, 15-21, 26-27.
A	PAN, G. et al. TRUNDD, A New Member of the TRAIL Receptor Family that Antagonizes TRAIL Signalling. FEBS Letters. 1998, Vol 424, pages 41-45.	1-13, 15-21, 26-27



Further documents are listed in the continuation of Box C.



See patent family annex.

\* Special categories of cited documents:

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\*

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\*

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\*

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

\*&amp;\*

document member of the same patent family

Date of the actual completion of the international search

03 AUGUST 1998

Date of mailing of the international search report

21 OCT 1998

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

  
NIRMAL S. BASU

Telephone No. (703) 308-0196

**INTERNATIONAL SEARCH REPORT**

International application No.  
PCT/US98/10981

**Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)**

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

**Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)**

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:  
1-13, 15-21, 23-24, 26-27

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US98/10981

### B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

APS, MEDLINE, JAPIO, BIOSIS, SCISEARCH, WPIDS, EMBASE, SPTREMB15, NGENSEQ31-2, EMBL-EST54, PIR56, SWISS-PROT25  
tr10 receptor, tumor necrosis factor, tm, apoptosis, tnfr-related apoptosis inducing ligand or trail, SEQ ID NO:1, 2, 7, 8 and 9

### BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Group I, claim(s) 1-13, 15-21, 23-24, 26-27, drawn to: a) nucleic acid encoding TR10 receptor (SEQ ID NO:1); b) polynucleotides that hybridize to said nucleic acid; c) recombinant vector and host cell for producing TR10 polypeptide; and d) method for treating a disease or disorder associated with inhibition of apoptosis.

Group II, claim 14, drawn to polynucleotide sequences of SEQ ID NO:7, 8 and 9.

Group III, claim 22, drawn to antibody that binds to a TR10 receptor.

Group IV, claim 25 drawn to a method of treating inflammatory disease and disorder by administering an antagonist of TR10.

The inventions listed as Groups I-IV do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons:

The main invention is Group I which is first product, first method of making the product, and first method of using the product. Pursuant to 37 CFR 1.474 (d), these claims are considered by the ISA/US to constitute the main invention, and none of the related groups II-IV correspond to the main invention.

The products of groups II and III, do not share the same or corresponding special technical feature with group I because they are drawn to products having materially different structures and functions, and each defines a separate invention over the art.

The methods of Group IV, do not share the same or corresponding special technical feature with group I because the methods of treatment use materially different compounds for materially different purposes, and each defines a separate invention over the art.

Therefore, the claims are so linked by a special technical feature within the meaning of PCT Rule 132.2 so as to form a single inventive concept.